KEAP1 Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	KEAP1, Polyclonal Antibody
Full Product Name	KEAP1 Antibody, FITC
Product Gene Name	anti-KEAP1 antibody
Product Synonym Names	Kelch-like ECH-associated protein 1; Cytosolic inhibitor of Nrf2; INrf2; Kelch-like protein 19; KEAP1; INRF2; KIAA0132; KLHL19
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation Purified
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.
Matching Pairs	Unconjugated Antibody: KEAP1 (MBS7003485) FITC Conjugated Antibody: KEAP1 (MBS1495835)
Molecular Weight	69,666 Da
OMIM	606016
3D Structure	ModBase 3D Structure for Q14145
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate	FITC
Immunogen	Recombinant human Kelch-like ECH-associated protein 1 protein
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-15244 / sc-15246 / sc-33894 / sc-33569
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-KEAP1 antibody	ELISA (EIA)
Related Product Information for anti-KEAP1 antibody	Acts as a substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1 and targets NFE2L2/NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. Retains NFE2L2/NRF2 and may also retain BPTF in the cytosol. Targets PGAM5 for ubiquitination and degradation by the proteasome.
Other Notes	Small volumes of anti-KEAP1 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for KEAP1 purchase	MBS1495835 is an antibody targeting KEAP1. Also known as KEAP1 Antibody, FITC. Reactive in Human. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for KEAP1. It may not necessarily be applicable to this product.

NCBI GI #	22027642
NCBI GeneID	9817
NCBI Accession #	NP_036421.2
UniProt Primary Accession #	Q14145
UniProt Secondary Accession #	Q6LEP0; Q8WTX1; Q9BPY9; B3KPD5
UniProt Related Accession #	Q14145
GenBank Nucleotide #	NM_012289.3
NCBI Official Full Name	kelch-like ECH-associated protein 1
NCBI Official Synonym Full Names	kelch like ECH associated protein 1
NCBI Official Symbol	KEAP1
NCBI Official Synonym Symbols	INrf2; KLHL19
UniProt Protein Name	Kelch-like ECH-associated protein 1
UniProt Synonym Protein Names	Cytosolic inhibitor of Nrf2; INrf2; Kelch-like protein 19
Protein Family	Kelch-like ECH-associated protein
UniProt Gene Name	KEAP1
UniProt Synonym Gene Names	INRF2; KIAA0132; KLHL19; INrf2
UniProt Entry Name	KEAP1_HUMAN
NCBI Protein Information	kelch-like ECH-associated protein 1
NCBI Summary for KEAP1	This gene encodes a protein containing KELCH-1 like domains, as well as a BTB/POZ domain. Kelch-like ECH-associated protein 1 interacts with NF-E2-related factor 2 in a redox-sensitive manner and the dissociation of the proteins in the cytoplasm is followed by transportation of NF-E2-related factor 2 to the nucleus. This interaction results in the expression of the catalytic subunit of gamma-glutamylcysteine synthetase. Two alternatively spliced transcript variants encoding the same isoform have been found for this gene. [provided by RefSeq, Jul 2008]

UniProt Comments for KEAP1	KEAP1: a substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1. Has tumor suppressor activity. Retains NRF2, BPTF and PGAM5 in the cytosol. Targets NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and of antioxidant response element-mediated detoxifying enzyme gene expression. Somatic mutations of KEAP1 or NRF2 commonly occur in solid cancers, irrespective of histological type, resulting in the constitutive transcription of cytoprotective genes. Genetic disruption of the KEAP1/CUL3 E3 ubiquitin ligase complex leads to the activation of the NF-kappaB pathway in lung cancer. Alternative activation of the cytoprotective NRF2 pathway exist in the absence of mutations in KEAP1 and HRF2: WTX, PALB2, SQSTM1, and DPP3 can competitively bind KEAP1. These proteins and others that bind KEAP1 contain an ¿ETGE¿ amino acid motif, which matches the KEAP1 interaction motif found in NRF2. Binding of these ¿ETGE¿-containing proteins can displace NRF2, freeing it to translocate to the nucleus and implement its cytoprotective transcriptional pathway. The DPP3 gene is frequently overexpressed in squamous cell lung carcinomas with high wt NRF2 activity. Its expression level significantly correlates with the response rate and progression-free survival of platinum-based first-line chemotherapy. May be a useful biomarker for predicting the chemosensitivity of patients with advanced-stage NSCLC. Broadly expressed, with highest levels in skeletal muscle. Ubiquitination and subsequent degradation of PGAM5 is inhibited by oxidative stress and sulforaphane. Protein type: Motility/polarity/chemotaxis; Endoplasmic reticulum; Ubiquitin conjugating system; Transcription regulation Chromosomal Location of Human Ortholog: 19p13.2 Cellular Component: actin filament; cell-cell adherens junction; cytoplasm; cytosol; endoplasmic reticulum; focal adhesion; microtubule organizing center; midbody; nucleoplasm Molecular Function: protein binding; transcription factor binding; ubiquitin-protein ligase activity Biological Process: alkanesulfonate metabolic process; cytoplasmic sequestering of transcription factor; flavonoid metabolic process; in utero embryonic development; malate metabolic process; negative regulation of transcription factor activity; positive regulation of proteasomal ubiquitin-dependent protein catabolic process; proteasomal ubiquitin-dependent protein catabolic process; proteasomal ubiquitin-independent protein catabolic process; protein oligomerization; protein ubiquitination; protein ubiquitination during ubiquitin-dependent protein catabolic process; regulation of epidermal cell differentiation; response to metal ion; selenium metabolic process; transcription, DNA-dependent
Research Articles on KEAP1	**1.** In the title.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/keap1/1495835