
Transferrin (TRF) Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	Transferrin (TRF), Polyclonal Antibody
Full Product Name	Biotin-Linked Antibody to Transferrin (TRF)
Product Gene Name	anti-TRF antibody
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, Pig
Form/Format	Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3,50% glycerol.
Purity/Purification	Affinity Chromatography
Concentration	500ug/ml (lot specific)
Specificity	The antibody is a rabbit polyclonal antibody raised against TRF. It has been selected for its ability to recognize TRF in immunohistochemical staining and western blotting.
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Matching Pairs	Unconjugated Antibody: TRF (MBS2001916) Biotin Conjugated Antibody: TRF (MBS2003784)
Matching Pairs	Biotin Conjugated Antibody: TRF (MBS2003784) Immunogen: TRF (MBS2012214)
Molecular Weight	77,064 Da
Chromosome Location	Chromosome: 3; NC_000003.11 (133464977..133497850). Location: 3q22.1
OMIM	190000
3D Structure	ModBase 3D Structure for P02787
Supply Chain Verification	Manufactured in a lab with traceable raw materials. Bulk orders can typically be prepared to the customer's specifications, please inquire.
Product Categories/Family for anti-TRF antibody	Biotin Antibody
ISO Certification	Manufactured in an ISO 9001:2015 and ISO 13485:2016 Certified Laboratory.
Applications Tested/Suitable for anti-TRF antibody	WB, ICC, IHC-P, IHC-F, ELISA
Application Notes for anti-TRF antibody	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.
Related Product Information for anti-TRF antibody	MBS2003784 is an antibody targeting TRF. Also known as Biotin-Linked Antibody to Transferrin (TRF). Reactive in Human, Pig. Host: Rabbit.

Other Notes	Small volumes of anti-TRF antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.



Western Blot of anti-TRF antibody

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for TRF. It may not necessarily be applicable to this product.

NCBI GI #	339453
NCBI GeneID	7018
NCBI Accession #	AAA61140.1
UniProt Primary Accession #	P02787
UniProt Secondary Accession #	O43890; Q1HBA5; Q9NQB8; Q9UHV0
UniProt Related Accession #	P02787
NCBI Official Full Name	transferrin
NCBI Official Synonym Full Names	transferrin
NCBI Official Symbol	TF
NCBI Official Synonym Symbols	TFQTL1; PRO1557; PRO2086
UniProt Protein Name	Serotransferrin
UniProt Synonym Protein Names	Beta-1 metal-binding globulin; Siderophilin
Protein Family	TBP-related factor
UniProt Gene Name	TF
UniProt Synonym Gene Names	Transferrin
UniProt Entry Name	TRFE_HUMAN
NCBI Protein Information	serotransferrin; siderophilin; beta-1 metal-binding globulin
NCBI Summary for TRF	This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum. [provided by RefSeq, Sep 2009]

UniProt Comments for TRF	TF: Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation. Defects in TF are the cause of atransferrinemia (ATRAF). Atransferrinemia is rare autosomal recessive disorder characterized by iron overload and hypochromic anemia. Belongs to the transferrin family. Protein type: Secreted; Secreted, signal peptide Chromosomal Location of Human Ortholog: 3q22.1 Cellular Component: extracellular space; cell surface; early endosome; cytoplasmic membrane-bound vesicle; extracellular region; coated pit; recycling endosome; endocytic vesicle; perinuclear region of cytoplasm; late endosome; apical plasma membrane; basal plasma membrane; endosome membrane; vesicle Molecular Function: protein binding; ferric iron binding; ubiquitin protein ligase binding Biological Process: platelet activation; retinal homeostasis; platelet degranulation; cellular iron ion homeostasis; transferrin transport; blood coagulation; transmembrane transport Disease: Atransferrinemia
Research Articles on TRF	**1.** Studies indicate that several genes have been linked to iron homeostasis, including transferrin (TF), iron regulatory protein 1 (ACO1) and transferrin receptor 2 (TFR2).
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-pig-antibody/transferrin-trf/2003784