Complement Component 3a (C3A) Antibody – Mouse Monoclonal

PRODUCT OVERVIEW

Product Name	Complement Component 3a (C3a), Monoclonal Antibody
Full Product Name	APC-Linked Monoclonal Antibody to Complement Component 3a (C3a)
Product Gene Name	anti-C3a antibody
Product Synonym Names	Complement Component 3a
Clonality	Monoclonal
Isotype	IgG1 Kappa
Clone Number	C18
Host	Mouse
Species Reactivity	Human
Form/Format	Liquid. 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Purity/Purification	Protein A + Protein G affinity chromatography
Concentration	>=0.1mg/mL (Please refer to the vial label for the specific concentration.) (lot specific)
Preparation and Storage	Store at 2 degree C to 8 degree C for frequent use, -20 degree C for 12 months Stable for 12 months if correctly stored.
Matching Pairs	Antibody: C3a Clone #C18 (MBS2147639) Immunogen: C3a (MBS2018643)
Matching Pairs	Unconjugated Antibody: C3a Clone #C18 (MBS2152375) APC Conjugated Antibody: C3a Clone #C18 (MBS2147639)
Molecular Weight	187,148 Da
Immunogen Sequence Positions	Ser672~Arg748
Chromosome Location	Chromosome: 19; NC_000019.9 (6677846..6720662, complement). Location: 19p13.3-p13.2
OMIM	120700
3D Structure	ModBase 3D Structure for P01024
Supply Chain Verification	Manufactured in a lab with traceable raw materials. Bulk orders can typically be prepared to the customer's specifications, please inquire.
Conjugation	APC
Product Categories/Family for anti-C3a antibody	Monoclonal Antibodies (Primary Abs); Primary, APC-conjugated; Infection immunity
ISO Certification	Manufactured in an ISO 9001:2015 and ISO 13485:2016 Certified Laboratory.
Applications Tested/Suitable for anti-C3a antibody	Western Blot (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)
Application Notes for anti-C3a antibody	WB: 0.5-2ug/mL IHC: 5-20ug/mL ICC: 5-20ug/mL

Related Product Information for anti-C3a antibody	MBS2147639 is an antibody targeting C3a. Also known as APC-Linked Monoclonal Antibody to Complement Component 3a (C3a). Reactive in Human. Host: Mouse.
Other Notes	Small volumes of anti-C3a antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for C3a. It may not necessarily be applicable to this product.

NCBI GI #	152012784
NCBI GeneID	718
NCBI Accession #	AAI50180.1
UniProt Primary Accession #	P01024
UniProt Secondary Accession #	A7E236
UniProt Related Accession #	P01024
NCBI Official Full Name	Complement component 3
NCBI Official Synonym Full Names	complement component 3
NCBI Official Symbol	C3
NCBI Official Synonym Symbols	ASP; C3a; C3b; AHUS5; ARMD9; CPAMD1
UniProt Protein Name	Complement C3
UniProt Synonym Protein Names	C3 and PZP-like alpha-2-macroglobulin domain-containing protein 1Cleaved into the following 11 chains:Complement C3 beta chainComplement C3 alpha chainC3a anaphylatoxinAcylation stimulating protein; ASP; Alternative name(s):; C3adesArg
UniProt Gene Name	C3
UniProt Synonym Gene Names	CPAMD1; ASP
UniProt Entry Name	CO3_HUMAN
NCBI Protein Information	complement C3; prepro-C3; C3a anaphylatoxin; complement component C3; complement component C3a; complement component C3b; acylation-stimulating protein cleavage product; C3 and PZP-like alpha-2-macroglobulin domain-containing protein 1
NCBI Summary for C3a	Complement component C3 plays a central role in the activation of complement system. Its activation is required for both classical and alternative complement activation pathways. People with C3 deficiency are susceptible to bacterial infection. [provided by RefSeq, Feb 2009]

UniProt Comments for C3a	C3: C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates. Defects in C3 are the cause of complement component 3 deficiency (C3D). A rare defect of the complement classical pathway. Patients develop recurrent, severe, pyogenic infections because of ineffective opsonization of pathogens. Some patients may also develop autoimmune disorders, such as arthralgia and vasculitic rashes, lupus-like syndrome and membranoproliferative glomerulonephritis. Genetic variation in C3 is associated with susceptibility to age-related macular degeneration type 9 (ARMD9). ARMD is a multifactorial eye disease and the most common cause of irreversible vision loss in the developed world. In most patients, the disease is manifest as ophthalmoscopically visible yellowish accumulations of protein and lipid that lie beneath the retinal pigment epithelium and within an elastin- containing structure known as Bruch membrane. Defects in C3 are a cause of susceptibility to hemolytic uremic syndrome atypical type 5 (AHUS5). An atypical form of hemolytic uremic syndrome. It is a complex genetic disease characterized by microangiopathic hemolytic anemia, thrombocytopenia, renal failure and absence of episodes of enterocolitis and diarrhea. In contrast to typical hemolytic uremic syndrome, atypical forms have a poorer prognosis, with higher death rates and frequent progression to end-stage renal disease. Susceptibility to the development of atypical hemolytic uremic syndrome can be conferred by mutations in various components of or regulatory factors in the complement cascade system. Other genes may play a role in modifying the phenotype. Increased levels of C3 and its cleavage product ASP, are associated with obesity, diabetes and coronary heart disease. Short-term endurance training reduces baseline ASP levels and subsequently fat storage. Protein type: Secreted, signal peptide; Secreted; Inhibitor Chromosomal Location of Human Ortholog: 19p13.3-p13.2 Cellular Component: extracellular space; plasma membrane; extracellular region Molecular Function: protein binding; endopeptidase inhibitor activity; C5L2 anaphylatoxin chemotactic receptor binding; receptor binding Biological Process: regulation of immune response; complement activation, alternative pathway; signal transduction; fatty acid metabolic process; complement activation; G-protein coupled receptor protein signaling pathway; positive regulation of angiogenesis; positive regulation of activation of membrane attack complex; positive regulation of G-protein coupled receptor protein signaling pathway; positive regulation of type IIa hypersensitivity; regulation of complement activation; innate immune response; immune response; positive regulation of protein amino acid phosphorylation; inflammatory response; complement activation, classical pathway Disease: Complement Component 3 Deficiency, Autosomal Recessive; Hemolytic Uremic Syndrome, Atypical, Susceptibility To, 5; Macular Degeneration, Age-related, 9
Research Articles on C3a	**1.** This study shows an association between a common single nucleotide polymorphism in C3 and susceptibility for community-acquired bacterial meningitis.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/monoclonal-human-antibody/complement-component-3a-c3a/2147639