Apolipoprotein E (APOE) Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	Apolipoprotein E (APOE), Polyclonal Antibody
Full Product Name	APC_Cy7-Linked Polyclonal Antibody to Apolipoprotein E (APOE)
Product Gene Name	anti-APOE antibody
Product Synonym Names	Apolipoprotein E; Apo-E; AD2; Apoprotein; Alzheimer Disease 2(E4-Associated,Late Onset
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Mouse
Form/Format	Liquid. 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Purity/Purification	Affinity Chromatography
Concentration	>=0.1mg/mL (Please refer to the vial label for the specific concentration.) (lot specific)
Preparation and Storage	Store at 2 degree C to 8 degree C for frequent use, -20 degree C for 12 months Stable for 12 months if correctly stored.
Matching Pairs	Antibody: APOE (MBS2150046) Immunogen: APOE (MBS2010968)
Matching Pairs	Unconjugated Antibody: APOE (MBS2151624) APC-CY7 Conjugated Antibody: APOE (MBS2150046)
Molecular Weight	35,867 Da
Immunogen Sequence Positions	Thr14~Lys167
3D Structure	ModBase 3D Structure for P08226
Supply Chain Verification	Manufactured in a lab with traceable raw materials. Bulk orders can typically be prepared to the customer's specifications, please inquire.
Conjugation	APC Cy7
Product Categories/Family for anti-APOE antibody	Polyclonal Antibodies (Primary Abs); Primary, APC_Cy7-conjugated; Metabolic pathway
ISO Certification	Manufactured in an ISO 9001:2015 and ISO 13485:2016 Certified Laboratory.
Applications Tested/Suitable for anti-APOE antibody	Western Blot (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)
Application Notes for anti-APOE antibody	WB: 0.5-2ug/mL IHC: 5-20ug/mL ICC: 5-20ug/mL
Related Product Information for anti-APOE antibody	MBS2150046 is an antibody targeting APOE. Also known as APC_Cy7-Linked Polyclonal Antibody to Apolipoprotein E (APOE). Reactive in Mouse. Host: Rabbit.
Other Notes	Small volumes of anti-APOE antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for APOE. It may not necessarily be applicable to this product.

NCBI GI #	802084029
NCBI GeneID	11816
NCBI Accession #	NP_001292748.1
UniProt Primary Accession #	P08226
UniProt Related Accession #	P08226
GenBank Nucleotide #	NM_001305819.1
NCBI Official Full Name	apolipoprotein E
NCBI Official Synonym Full Names	apolipoprotein E
NCBI Official Symbol	Apoe
NCBI Official Synonym Symbols	Apo-E; AI255918
UniProt Protein Name	Apolipoprotein E
Protein Family	Apolipoprotein
UniProt Gene Name	Apoe
UniProt Synonym Gene Names	Apo-E
UniProt Entry Name	APOE_MOUSE
NCBI Protein Information	apolipoprotein E
NCBI Summary for APOE	This gene encodes a member of the apolipoprotein A1/A4/E family of proteins. This protein is involved in the transport of lipoproteins in the blood. It binds to a specific liver and peripheral cell receptor, and is essential for the normal catabolism of triglyceride-rich lipoprotein constituents. Homozygous knockout mice for this gene accumulate high levels of cholesterol in the blood and develop atherosclerosis. Different alleles of this gene have been associated with either increased risk or a protective effect for Alzheimer's disease in human patients. This gene maps to chromosome 7 in a cluster with the related apolipoprotein C1, C2 and C4 genes. [provided by RefSeq, Apr 2015]

UniProt Comments for APOE	APOE: Mediates the binding, internalization, and catabolism of lipoprotein particles. It can serve as a ligand for the LDL (apo B/E) receptor and for the specific apo-E receptor (chylomicron remnant) of hepatic tissues. Defects in APOE are a cause of hyperlipoproteinemia type 3 (HLPP3); also known as familial dysbetalipoproteinemia. Individuals with HLPP3 are clinically characterized by xanthomas, yellowish lipid deposits in the palmar crease, or less specific on tendons and on elbows. The disorder rarely manifests before the third decade in men. In women, it is usually expressed only after the menopause. The vast majority of the patients are homozygous for APOE*2 alleles. More severe cases of HLPP3 have also been observed in individuals heterozygous for rare APOE variants. The influence of APOE on lipid levels is often suggested to have major implications for the risk of coronary artery disease (CAD). Individuals carrying the common APOE*4 variant are at higher risk of CAD. Genetic variations in APOE are associated with Alzheimer disease type 2 (AD2). It is a late-onset neurodegenerative disorder characterized by progressive dementia, loss of cognitive abilities, and deposition of fibrillar amyloid proteins as intraneuronal neurofibrillary tangles, extracellular amyloid plaques and vascular amyloid deposits. The major constituent of these plaques is the neurotoxic amyloid-beta-APP 40-42 peptide (s), derived proteolytically from the transmembrane precursor protein APP by sequential secretase processing. The cytotoxic C-terminal fragments (CTFs) and the caspase-cleaved products such as C31 derived from APP, are also implicated in neuronal death. The APOE*4 allele is genetically associated with the common late onset familial and sporadic forms of Alzheimer disease. Risk for AD increased from 20% to 90% and mean age at onset decreased from 84 to 68 years with increasing number of APOE*4 alleles in 42 families with late onset AD. Thus APOE*4 gene dose is a major risk factor for late onset AD and, in these families, homozygosity for APOE*4 was virtually sufficient to cause AD by age 80. The mechanism by which APOE*4 participates in pathogenesis is not known. Defects in APOE are a cause of sea-blue histiocyte disease (SBHD); also known as sea-blue histiocytosis. This disorder is characterized by splenomegaly, mild thrombocytopenia and, in the bone marrow, numerous histiocytes containing cytoplasmic granules which stain bright blue with the usual hematologic stains. The syndrome is the consequence of an inherited metabolic defect analogous to Gaucher disease and other sphingolipidoses. Defects in APOE are a cause of lipoprotein glomerulopathy (LPG). LPG is an uncommon kidney disease characterized by proteinuria, progressive kidney failure, and distinctive lipoprotein thrombi in glomerular capillaries. It mainly affects people of Japanese and Chinese origin. The disorder has rarely been described in Caucasians. Belongs to the apolipoprotein A1/A4/E family. Protein type: Lipid-binding; Secreted; Secreted, signal peptide Cellular Component: cell soma; cell surface; chylomicron; cytoplasm; dendrite; early endosome; endoplasmic reticulum; endosome; extracellular matrix; extracellular space; extrinsic to external side of plasma membrane; Golgi apparatus; late endosome; lysosome; membrane; microtubule; nuclear envelope; nucleus; plasma membrane Molecular Function: antioxidant activity; beta-amyloid binding; cholesterol binding; cholesterol transporter activity; heparin binding; hydroxyapatite binding; identical protein binding; lipid binding; lipid transporter activity; low-density lipoprotein receptor binding; metal chelating activity; phospholipid binding; protein binding; protein homodimerization activity; receptor binding; tau protein binding Biological Process: aging; artery morphogenesis; axon regeneration; cellular calcium ion homeostasis; cGMP-mediated signaling; cholesterol biosynthetic process; cholesterol catabolic process; cholesterol efflux; cholesterol homeostasis; cholesterol metabolic process; fatty acid homeostasis; G-protein coupled receptor protein signaling pathway; lipid homeostasis; lipid metabolic process; lipid transport; lipoprotein biosynthetic process; lipoprotein catabolic process; lipoprotein metabolic process; long-chain fatty acid transport; maintenance of cellular localization; negative regulation of blood coagulation; negative regulation of blood vessel endothelial cell migration; negative regulation of cholesterol biosynthetic process; negative regulation of endothelial cell proliferation; negative regulation of inflammatory response; negative regulation of lipid biosynthetic process; negative regulation of MAP kinase activity; negative regulation of neuron apoptosis; nitric oxide mediated signal transduction; phospholipid efflux; positive regulation of axon extension; positive regulation of cGMP biosynthetic process; positive regulation of lipid biosynthetic process; positive regulation of low-density lipoprotein receptor catabolic process; positive regulation of membrane protein ectodomain proteolysis; positive regulation of nitric-oxide synthase activity; protein import; protein localization; receptor-mediated endocytosis; regulation of axon extension; regulation of Cdc42 protein signal transduction; regulation of cholesterol transport; regulation of gene expression; response to dietary excess; response to oxidative stress; reverse cholesterol transport; triacylglycerol catabolic process; triacylglycerol metabolic process; vasodilation; virus assembly
Research Articles on APOE	**1.** Sex Difference in Coronary Endothelial Dysfunction in Apolipoprotein E Knockout Mouse: Role of NO and A2A Adenosine Receptor.

Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-mouse-antibody/apolipoprotein-e-apoe/2150046