
TLR3 Antibody – Rabbit Polyclonal


PRODUCT OVERVIEW

Product Name	TLR3, Polyclonal Antibody
Full Product Name	TLR3 Polyclonal Antibody
Product Gene Name	anti-TLR3 antibody
Product Synonym Names	TLR3; CD283; IIAE2
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human, Mouse, Rat
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Purity/Purification	Affinity purification
Concentration	1mg/mL (lot specific)
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.
Molecular Weight	103,829 Da
Chromosome Location	Chromosome: 4; NC_000004.11 (186990309..187006252). Location: 4q35
OMIM	603029
3D Structure	ModBase 3D Structure for O15455
Immunogen	Recombinant fusion protein of human TLR3 (NP_0032561)
Conjugation	Unconjugated
Product Categories/Family for anti-TLR3 antibody	Cancer; Immunology; Microbiology; Signal Transduction; Polyclonal Antibody
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-TLR3 antibody	Immunofluorescence (IF)
Application Notes for anti-TLR3 antibody	IF: 1:50-1:200
Related Product Information for anti-TLR3 antibody	The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of the leukocytes. It recognizes dsRNA associated with viral infection, and induces the activation of NF-kappaB and the production of type I interferons. It may thus play a role in host defense against viruses. Use of alternative polyadenylation sites to generate different length transcripts has been noted for this gene.
Other Notes	Small volumes of anti-TLR3 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.


Searchable Terms for TLR3 purchase	MBS2564672 is an antibody targeting TLR3. Also known as TLR3 Polyclonal Antibody. Reactive in Human, Mouse, Rat. Host: Rabbit.



Immunofluorescence (IF) of anti-TLR3 antibody
Immunofluorescence analysis of C6 cells using TLR3 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence (IF) of anti-TLR3 antibody
Immunofluorescence analysis of HeLa cells using TLR3 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence (IF) of anti-TLR3 antibody
Immunofluorescence analysis of L929 cells using TLR3 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for TLR3. It may not necessarily be applicable to this product.

NCBI GI #	86161330
NCBI GeneID	7098
NCBI Accession #	ABC86910.1
UniProt Primary Accession #	O15455
UniProt Secondary Accession #	Q4VAL2
UniProt Related Accession #	O15455
NCBI Official Full Name	TLR3
NCBI Official Synonym Full Names	toll-like receptor 3
NCBI Official Symbol	TLR3
NCBI Official Synonym Symbols	CD283; IIAE2
UniProt Protein Name	Toll-like receptor 3
Protein Family	Toll-like receptor

UniProt Gene Name	TLR3
UniProt Entry Name	TLR3_HUMAN
NCBI Protein Information	toll-like receptor 3
UniProt Comments for TLR3	TLR3: a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. Most abundantly expressed in placenta and pancreas. Restricted to the dendritic subpopulation of the leukocytes. It recognizes dsRNA associated with viral infection. Acts via MyD88 and TRAF6, leading to the activation of NF-kappaB and the production of type I interferon. Protein type: Receptor, cytokine; Membrane protein, integral Chromosomal Location of Human Ortholog: 4q35 Cellular Component: Golgi membrane; endoplasmic reticulum membrane; cell surface; membrane; lysosomal membrane; integral to plasma membrane; cytoplasm; endosome membrane Molecular Function: identical protein binding; protein binding; transmembrane receptor activity; double-stranded RNA binding; receptor activity Biological Process: negative regulation of osteoclast differentiation; I-kappaB kinase/NF-kappaB cascade; positive regulation of interleukin-12 production; positive regulation of apoptosis; positive regulation of JNK cascade; positive regulation of NF-kappaB import into nucleus; positive regulation of interferon-alpha biosynthetic process; toll-like receptor 3 signaling pathway; signal transduction; activation of NF-kappaB transcription factor; positive regulation of interleukin-8 production; response to exogenous dsRNA; positive regulation of interferon-beta production; inflammatory response; defense response to virus; toll-like receptor 4 signaling pathway; MyD88-independent toll-like receptor signaling pathway; male gonad development; activation of NF-kappaB-inducing kinase; positive regulation of interleukin-6 production; positive regulation of tumor necrosis factor production; positive regulation of chemokine production; positive regulation of toll-like receptor signaling pathway; hyperosmotic response; positive regulation of chemokine biosynthetic process; detection of virus; defense response to bacterium; positive regulation of interferon-beta biosynthetic process; toll-like receptor signaling pathway; innate immune response; positive regulation of transcription from RNA polymerase II promoter; I-kappaB phosphorylation; microglial cell activation during immune response; positive regulation of inflammatory response; positive regulation of interferon-gamma biosynthetic process Disease: Human Immunodeficiency Virus Type 1, Susceptibility To
Research Articles on TLR3	**1.** TLR3 +1234C/T polymorphism is associated with HBV-related hepatocellular carcinoma.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.