AMPKalpha2 Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	AMPKalpha2, Polyclonal Antibody
Full Product Name	AMPKalpha2 Polyclonal Antibody
Product Gene Name	anti-AMPKalpha2 antibody
Product Synonym Names	PRKAA2; AMPK; AMPK2; AMPKa2; PRKAA
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human, Rat
Form/Format	PBS with 0.05% proclin300,pH7.3.
Purity/Purification	Affinity purification
Concentration	1mg/mL (lot specific)
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.
Molecular Weight	62,320 Da
OMIM	600497
3D Structure	ModBase 3D Structure for P54646
Immunogen	Recombinant fusion protein of human PRKAA2
Research Area	Cancer, Cardiovascular, Metabolism, Signal Transduction
Product Categories/Family for anti-AMPKalpha2 antibody	Antibodies; Primary Antibodies; Polyclonal Antibody
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-AMPKalpha2 antibody	Western Blot (WB), Immunofluorescence (IF)
Application Notes for anti-AMPKalpha2 antibody	WB: 1:500-1:1000 IF: 1:50-1:200
Related Product Information for anti-AMPKalpha2 antibody	The protein encoded by this gene is catalytic subunit of the AMP-activated protein kinase (AMPK). AMPK is heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. Studies of the mouse counterpartuggest that this catalytic subunit may control whole-body insulin sensitivity and is necessary for maintaining myocardial energy homeostasis during ischemia.
Other Notes	Small volumes of anti-AMPKalpha2 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for AMPKalpha2 purchase	MBS2572718 is an antibody targeting AMPKalpha2. Also known as AMPKalpha2 Polyclonal Antibody. Reactive in Human, Rat. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for AMPKalpha2. It may not necessarily be applicable to this product.

NCBI GI #	786491
NCBI GeneID	5563
NCBI Accession #	AAB32732.1
UniProt Primary Accession #	P54646
UniProt Secondary Accession #	Q9H1E8; Q9UD43
UniProt Related Accession #	P54646; AAB32732
NCBI Official Full Name	AMP-activated protein kinase, AMPK
NCBI Official Synonym Full Names	protein kinase, AMP-activated, alpha 2 catalytic subunit
NCBI Official Symbol	PRKAA2
NCBI Official Synonym Symbols	AMPK; AMPK2; PRKAA; AMPKa2
UniProt Protein Name	5'-AMP-activated protein kinase catalytic subunit alpha-2
UniProt Synonym Protein Names	Acetyl-CoA carboxylase kinase (EC:2.7.11.27); ACACA kinase; Hydroxymethylglutaryl-CoA reductase kinase (EC:2.7.11.31); HMGCR kinase
UniProt Gene Name	PRKAA2
UniProt Synonym Gene Names	AMPK; AMPK2; AMPK subunit alpha-2; ACACA kinase; HMGCR kinase
UniProt Entry Name	AAPK2_HUMAN
NCBI Protein Information	5'-AMP-activated protein kinase catalytic subunit alpha-2; ACACA kinase; HMGCR kinase; AMPK-alpha-2 chain; AMPK subunit alpha-2; acetyl-CoA carboxylase kinase; hydroxymethylglutaryl-CoA reductase kinase; AMP-activated protein kinase alpha-2 subunit varian
NCBI Summary for AMPKalpha2	The protein encoded by this gene is a catalytic subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. Studies of the mouse counterpart suggest that this catalytic subunit may control whole-body insulin sensitivity and is necessary for maintaining myocardial energy homeostasis during ischemia. [provided by RefSeq, Jul 2008]

UniProt Comments for AMPKalpha2	AMPKA2: a catalytic subunit of AMP-activated protein kinase (AMPK). Acts as an energy sensor, playing a key role in regulating cellular energy metabolism. A protein kinase of the CAMKL family whose activation is regulated by the balance between ADP/AMP/ATP, and intracellular Ca(2+) levels. Acts as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-ADP and -AMP rise in response to fuel limitation and/or hypoxia. Activates energy-producing pathways and inhibits energy-consuming processes. Restores ATP levels in cells by switching off anabolic and switching on catabolic pathways. Activated primarily by rising ADP levels and not, as previously thought, solely by AMP. AMPK resembles an adenylate charge regulatory system in which anabolic and catabolic pathways are regulated by adenine nucleotide ratios. Acts via direct phosphorylation of metabolic enzymes and transcription regulators. Regulates fatty acid synthesis by phosphorylating acetyl-CoA carboxylase. Regulates cholesterol synthesis by phosphorylating and inactivating hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Activated by at least two distinct upstream kinases: the tumor suppressor LKB1 and CaMKK2. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton, probably by indirectly activating myosin. AMPK is a heterotrimer of an alpha catalytic subunit (AMPKA1 or -2), a beta (AMPKB1 or -2) and a gamma non-catalytic subunit (AMPKG1, -2 or -3). Different possible combinations of subunits give rise to 12 different holoenzymes. Binding of ADP or AMP to non-catalytic gamma subunit (PRKAG1, -2 or -3) results in allosteric activation. AMPK is activated by antihyperglycemic drug metformin, a drug prescribed to patients with type 2 diabetes: in vivo, metformin seems to mainly inhibit liver gluconeogenesis. However, metformin can be used to activate AMPK in muscle and other cells in culture or ex vivo. Selectively inhibited by compound C (6-[4-(2-Piperidin-1-yl-ethoxy)-phenyl)]-3-pyridin-4-yl-pyyrazolo[1,5-a] pyrimidine. Activated by resveratrol, a natural polyphenol present in red wine, and S17834, a synthetic polyphenol. Salicylate/aspirin directly activates kinase activity. Studies in the mouse suggest that AMPK2 may control whole-body insulin sensitivity and is necessary for maintaining myocardial energy homeostasis during ischemia. Protein type: Protein kinase, Ser/Thr (non-receptor); EC 2.7.11.27; EC 2.7.11.1; Kinase, protein; Autophagy; Protein kinase, CAMK; EC 2.7.11.31; CAMK group; CAMKL family; AMPK subfamily Chromosomal Location of Human Ortholog: 1p31 Cellular Component: nucleoplasm; cytosol Molecular Function: AMP-activated protein kinase activity; protein serine/threonine kinase activity; protein binding; metal ion binding; [acetyl-CoA carboxylase] kinase activity; protein serine/threonine/tyrosine kinase activity; chromatin binding; histone serine kinase activity; ATP binding; protein kinase activity; [hydroxymethylglutaryl-CoA reductase (NADPH)] kinase activity Biological Process: lipid biosynthetic process; rhythmic process; cellular lipid metabolic process; carnitine shuttle; glucose homeostasis; signal transduction; protein amino acid phosphorylation; cellular response to glucose starvation; cellular response to nutrient levels; regulation of fatty acid biosynthetic process; regulation of transcription, DNA-dependent; response to stress; cell cycle arrest; positive regulation of autophagy; fatty acid biosynthetic process; mitochondrion organization and biogenesis; negative regulation of TOR signaling pathway; Wnt receptor signaling pathway; transcription, DNA-dependent; organelle organization and biogenesis; regulation of circadian rhythm; cholesterol biosynthetic process; fatty acid homeostasis; positive regulation of glycolysis; insulin receptor signaling pathway; energy reserve metabolic process; autophagy; negative regulation of apoptosis
Research Articles on AMPKalpha2	**1.** High PRKAA2 expression is associated with follicular lymphoma.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.

Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-rat-antibody/ampkalpha2/2572718