APP Antibody – Goat Polyclonal

PRODUCT OVERVIEW

Product Name	APP, Polyclonal Antibody
Full Product Name	APP antibody
Product Gene Name	anti-APP antibody
Product Synonym Names	Polyclonal APP; Anti-APP; Amyloid Precursor Protein
Clonality	Polyclonal
Host	Goat
Species Reactivity	Human
Form/Format	PBS, pH 7.4 with 0.05% NaN3.
Specificity	Human
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.
Molecular Weight	84,521 Da
OMIM	104300
Biohazard Information	This product contains sodium azide as preservative. Although the amount of sodium azide is very small appropriate care must be taken when handling.
Immunogen	APP antibody was raised in goat using a synthetic peptide corresponding to amino acids 44-63 of the N-terminus of the alzheimer's disease APP as the immunogen.
Product Categories/Family for anti-APP antibody	Neuroscience
Applications Tested/Suitable for anti-APP antibody	ELISA (EIA)
Application Notes for anti-APP antibody	ELISA: 1:5,000
Related Product Information for anti-APP antibody	MBS535989 is an antibody targeting APP. Also known as APP antibody. Reactive in Human. Host: Goat. Goat polyclonal APP antibody
Other Notes	Small volumes of anti-APP antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for APP. It may not necessarily be applicable to this product.

NCBI GI #	236721
NCBI GeneID	351
NCBI Accession #	AAB19991.1
UniProt Secondary Accession #	P09000; P78438; Q13764; Q13778; Q13793; Q16011; B2R5V1; B4DII8; D3DSD1; D3DSD2; D3DSD3
UniProt Related Accession #	P05067

NCBI Official Full Name	amyloid protein, partial
NCBI Official Synonym Full Names	amyloid beta (A4) precursor protein
NCBI Official Symbol	APP
NCBI Official Synonym Symbols	AAA; AD1; PN2; ABPP; APPI; CVAP; ABETA; PN-II; CTFgamma
UniProt Protein Name	Amyloid beta A4 protein
UniProt Synonym Protein Names	ABPP; APPI; APP
Protein Family	Amyloid beta A4 protein
UniProt Gene Name	APP
UniProt Synonym Gene Names	A4; AD1; APP; CVAP; PN-II; S-APP-alpha; S-APP-beta; AICD-59; AID(59); AICD-57; AID(57); AICD-50; AID(50)
UniProt Entry Name	A4_HUMAN
NCBI Protein Information	amyloid beta A4 protein
NCBI Summary for APP	This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. In addition, two of the peptides are antimicrobial peptides, having been shown to have bacteriocidal and antifungal activities. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene. [provided by RefSeq, Aug 2014]

UniProt Comments for APP	APP: a cell surface receptor that influences neurite growth, neuronal adhesion and axonogenesis. Cleaved by secretases to form a number of peptides, some of which bind to the acetyltransferase complex Fe65/TIP60 to promote transcriptional activation. The Abeta peptide is released from the cell, its extracellular deposition and accumulation form the main components of amyloid plaques in Alzheimer's disease. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis. Can promote transcription activation through binding to Fe65-Tip60 and inhibits Notch signaling through interaction with Numb. Couples to apoptosis-inducing pathways such as those mediated by G(O) and JIP. Inhibits G(O) alpha ATPase activity. Acts as a kinesin I membrane receptor, mediating the axonal transport of beta-secretase and presenilin 1. Involved in copper homeostasis/oxidative stress through copper ion reduction. In vitro, copper-metallated APP induces neuronal death directly or is potentiated through Cu(2+)-mediated low-density lipoprotein oxidation. Can regulate neurite outgrowth through binding to components of the extracellular matrix such as heparin and collagen I and IV. Induces a RAGE-dependent pathway that activates p38 MAPK, resulting in internalization of amyloid-beta peptide and leading to mitochondrial dysfunction in cultured cortical neurons. Provides Cu(2+) ions for GPC1 which are required for release of nitric oxide (NO) and subsequent degradation of the heparan sulfate chains on GPC1. Binds, via its C-terminus, to the PID domain of several cytoplasmic proteins, including APBB family members, the APBA family, JIP1, SHC1 and, NUMB and DAB1. Binding to DAB1 inhibits its serine phosphorylation. Associates with microtubules in the presence of ATP and in a kinesin-dependent manner. Amyloid beta-42 binds nAChRA7 in hippocampal neurons. Beta-amyloid associates with HADH2. Soluble APP binds, via its N-terminal head, to FBLN1. Expressed in all fetal tissues examined with highest levels in brain, kidney, heart and spleen. Weak expression in liver. In adult brain, highest expression found in the frontal lobe of the cortex and in the anterior perisylvian cortex-opercular gyri. Moderate expression in the cerebellar cortex, the posterior perisylvian cortex-opercular gyri and the temporal associated cortex. Weak expression found in the striate, extra- striate and motor cortices. Expressed in cerebrospinal fluid, and plasma. 10 isoforms of the human protein are produced by alternative splicing. Isoform APP695 is the predominant form in neuronal tissue, isoform APP751 and isoform APP770 are widely expressed in non- neuronal cells. Isoform APP751 is the most abundant form in T-lymphocytes. Appican is expressed in astrocytes. The splice isoforms that contain the BPTI domain possess protease inhibitor activity. Belongs to the APP family. Protein type: Apoptosis; Membrane protein, integral; Cell surface; Receptor, misc.; Transcription factor Chromosomal Location of Human Ortholog: 21q21.3 Cellular Component: Golgi apparatus; extracellular space; cell surface; integral to plasma membrane; integral to membrane; coated pit; intercellular junction; cytosol; ER to Golgi transport vesicle; lipid raft; ciliary rootlet; perinuclear region of cytoplasm; nuclear envelope lumen; cytoplasm; synapse; dendritic shaft; neuromuscular junction; receptor complex; endosome; intracellular membrane-bound organelle; dendritic spine; extracellular region; apical part of cell; axon; plasma membrane; spindle midzone Molecular Function: heparin binding; serine-type endopeptidase inhibitor activity; identical protein binding; protein binding; protease activator activity; enzyme binding; DNA binding; transition metal ion binding; PTB domain binding; acetylcholine receptor binding; receptor binding Biological Process: extracellular matrix organization and biogenesis; adult locomotory behavior; locomotory behavior; mRNA polyadenylation; positive regulation of mitotic cell cycle; protein amino acid phosphorylation; regulation of translation; platelet degranulation; synaptic growth at neuromuscular junction; forebrain development; dendrite development; visual learning; collateral sprouting in the absence of injury; neuromuscular process controlling balance; cell adhesion; neurite development; cholesterol metabolic process; platelet activation; Notch signaling pathway; cellular copper ion homeostasis; regulation of epidermal growth factor receptor activity; mating behavior; axon cargo transport; regulation of multicellular organism growth; endocytosis; axon midline choice point recognition; smooth endoplasmic reticulum calcium ion homeostasis; negative regulation of neuron differentiation; neuron apoptosis; suckling behavior; axonogenesis; ionotropic glutamate receptor signaling pathway; regulation of protein binding; regulation of synapse structure and activity; innate immune response; positive regulation of transcription from RNA polymerase II promoter; response to oxidative stress; blood coagulation; neuron remodeling Disease: Alzheimer Disease
Research Articles on APP	**1.** Abeta25-35 can induce the release from human neutrophils of pro-MMP-9, a metalloprotease involved in the onset of inflammation
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.

Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/app/535989