
HSF4 Antibody – Mouse Monoclonal

PRODUCT OVERVIEW

Product Name	HSF4, Monoclonal Antibody
Full Product Name	HSF4 (Heat Shock Transcription Factor 4, CTM) (HRP)
Product Gene Name	anti-HSF4 antibody
Product Synonym Names	Heat Shock Transcription Factor 4; CTM
Clonality	Monoclonal
Isotype	IgG2a,k
Clone Number	1A4
Host	Mouse
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with horseradish peroxidase (HRP).
Purity/Purification	Purified
Specificity	Recognizes HSF4.
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and stor at -20 degree C. Aliquots are stable at -20 degree C for 12 months after receipt. Dilute required amount only prior to immediate use. Further dilutions ca be made in assay buffer. Note: Sodium azide is a potent inhibitor of peroxidase and should not be added to HRP conjugates. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.
Molecular Weight	50kDa
OMIM	phenotype 602438
3D Structure	ModBase 3D Structure for Q9ULV5
Immunogen	HSF4 (NP_001529, 121aa-220aa) partial recombinant protein with GST tag. MW of the GST tag alone is 26kD.
Immunogen Sequence	KVPALRGDDGRWRPEDLGRLLGEVQALRGVQESTEARLRELRQQNEILWREVVTLRQSHGQQHRVIGKLIQCLFGPLQAGPSNAGGKRKLSLMLDEGS
Conjugate	HRP
Product Categories/Family for anti-HSF4 antibody	Antibodies; Transcription Factors
Applications Tested/Suitable for anti-HSF4 antibody	Western Blot (WB)
Application Notes for anti-HSF4 antibody	Applications are based on unconjugated antibody.

Related Product Information for anti-HSF4 antibody	MBS6181120 is an antibody targeting HSF4. Also known as HSF4 (Heat Shock Transcription Factor 4, CTM) (HRP). Host: Mouse.
Other Notes	Small volumes of anti-HSF4 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.



Testing Data of anti-HSF4 antibody
Detection limit for recombinant GST tagged HSF4 is 0.3 ng/ml as a capture antibody.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for HSF4. It may not necessarily be applicable to this product.

NCBI GI #	100913211
NCBI GeneID	3299
NCBI Accession #	NP_001529
UniProt Primary Accession #	Q9ULV5
UniProt Related Accession #	Q9ULV5
GenBank Nucleotide #	NM_001538
NCBI Official Full Name	heat shock factor protein 4 isoform a
NCBI Official Synonym Full Names	heat shock transcription factor 4
NCBI Official Symbol	HSF4
NCBI Official Synonym Symbols	CTM; CTRCT5
UniProt Protein Name	Heat shock factor protein 4
UniProt Synonym Protein Names	Heat shock transcription factor 4; HSTF 4
Protein Family	Heat shock factor protein
UniProt Gene Name	HSF4
UniProt Synonym Gene Names	HSF 4; hHSF4; HSTF 4
UniProt Entry Name	HSF4_HUMAN
NCBI Protein Information	heat shock factor protein 4
NCBI Summary for HSF4	Heat-shock transcription factors (HSFs) activate heat-shock response genes under conditions of heat or other stresses. HSF4 lacks the carboxyl-terminal hydrophobic repeat which is shared among all vertebrate HSFs and has been suggested to be involved in the negative regulation of DNA binding activity. Two alternatively spliced transcripts encoding distinct isoforms and possessing different transcriptional activity have been described. [provided by RefSeq, Jul 2008]

UniProt Comments for HSF4	HSF4: DNA-binding protein that specifically binds heat shock promoter elements (HSE). Isoform HSF4A represses transcription while the isoform HSF4B activates transcription. Defects in HSF4 are the cause of cataract, zonular (CZ). A form of zonular cataract. Zonular or lamellar cataracts are opacities, broad or narrow, usually consisting of powdery white dots affecting only certain layers or zones between the cortex and nucleus of an otherwise clear lens. The opacity may be so dense as to render the entire central region of the lens completely opaque, or so translucent that vision is hardly if at all impeded. Zonular cataracts generally do not involve the embryonic nucleus, though sometimes they involve the fetal nucleus. Usually sharply separated from a clear cortex outside them, they may have projections from their outer edges known as riders or spokes. Defects in HSF4 are the cause of cataract Marner type (CAM). A form of cataract with variable and progressive opacities. Affected individuals present with zonular cataract, although some have nuclear, anterior polar, or stellate cataract. Finger malformation is observed in some kindreds. Belongs to the HSF family. 2 isoforms of the human protein are produced by alternative splicing. Protein type: Transcription, coactivator/corepressor; DNA-binding; Cell development/differentiation Chromosomal Location of Human Ortholog: 16q21 Cellular Component: nucleoplasm Molecular Function: sequence-specific DNA binding; transcription factor activity; transcription corepressor activity; protein phosphatase binding Biological Process: histone H3-K9 demethylation; camera-type eye development; transcription, DNA-dependent; regulation of transcription, DNA-dependent; positive regulation of cell proliferation; positive regulation of transcription from RNA polymerase II promoter; negative regulation of transcription from RNA polymerase II promoter; positive regulation of cell differentiation; cell development Disease: Cataract 5, Multiple Types
Research Articles on HSF4	**1.** Ectopic UAP56 upregulated HSF4-controlled HSP25 and alpha B-crystallin proteins expression, while knocking down UAP56 by shRNA reversed it.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/monoclonal-antibody/hsf4/6181120