
RET Antibody – Rabbit Polyclonal


PRODUCT OVERVIEW

Product Name	RET, Polyclonal Antibody
Full Product Name	RET, NT (Q28) (RET, CDHF12, CDHR16, PTC, RET51, Proto-oncogene tyrosine-protein kinase receptor Ret, Cadherin family member 12, Proto-oncogene c-Ret, Soluble RET kinase fragment, Extracellular cell-membrane anchored RET cadherin 120kD fragment) (MaxLight
Product Gene Name	anti-RET antibody
Product Synonym Names	RET; CDHF12; CDHR16; PTC; RET51; Proto-oncogene tyrosine-protein kinase receptor Ret; Cadherin family member 12; Proto-oncogene c-Ret; Soluble RET kinase fragment; Extracellular cell-membrane anchored RET cadherin 120kD fragment
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight405.
Purity/Purification	Purified by Protein A/G Affinity Chromatography.
Specificity	Recognizes human RET.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight405 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	124,319 Da
Chromosome Location	Chromosome: 10; NC_000010.10 (43572517..43625799). Location: 10q11.2
OMIM	114500
3D Structure	ModBase 3D Structure for P07949
Immunogen	KLH-conjugated synthetic peptide mapping to a fragment of residues within amino acids 13-44 in the N-terminal region of human RET.
Conjugate	MaxLight405
Note	Preservative Free
Product Categories/Family for anti-RET antibody	Antibodies; Protein Kinases
Applications Tested/Suitable for anti-RET antibody	Western Blot (WB), Flow Cytometry (FC/FACS), FLISA
Application Notes for anti-RET antibody	Applications are based on unconjugated antibody.
Related Product Information for anti-RET antibody	RET, a member of the cadherin superfamily, is one of the receptor tyrosine kinases, which are cell-surface molecules that transduce signals for cell growth and differentiation. This protein plays a crucial role in neural crest development, and the gene can undergo oncogenic activation in vivo and in vitro by cytogenetic rearrangement. Mutations i are associated with the disorders multiple endocrine neoplasia, type IIA, multiple endocrine neoplasia, type IIB, Hirschsprung disease, and medullary thyroid carcinoma.

Product References and Citations for anti-RET antibody	Da Silva, A.M., et al., J. Clin. Endocrinol. Metab. 88(11):5438-5443 (2003). McWhinney, S.R., et al., J. Clin. Endocrinol. Metab. 88(10):4911-4916 (2003). D'Alessio, A., et al., Endocrinology 144(10):4298-4305 (2003). Soares, P., et al., Oncogene 22(29):4578-4580 (2003). Punales, M.K., et al., J. Clin. Endocrinol. Metab. 88(6):2644-2649 (2003).
Other Notes	Small volumes of anti-RET antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for RET purchase	MBS6341394 is an antibody targeting RET. Also known as RET, NT (Q28) (RET, CDHF12, CDHR16, PTC, RET51, Proto-oncogene tyrosine-protein kinase receptor Ret, Cadherin family member 12, Proto-oncogene c-Ret, Soluble RET kinase fragment, Extracellular cell-membrane anchored RET cadherin 120kD fragment) (MaxLight. Reactive in Human. Host: Rabbit.



Western Blot (WB) of anti-RET antibody
Western Blot analysis of hRET-G28 in MCF7 cell line lysates (35ug/lane) using 40968. RET (arrow) was detected using the purified Pab.



Flow Cytometry (FC/FACS) of anti-RET antibody
Flow cytometric analysis of MCF-7 cells using 040968 (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for RET. It may not necessarily be applicable to this product.

NCBI GI #	10862701
NCBI GeneID	5979
NCBI Accession #	NP_065681.1
UniProt Primary Accession #	P07949
UniProt Secondary Accession #	Q15250; Q9BTB0; Q9H4A2; A8K6Z2
UniProt Related Accession #	P07949
GenBank Nucleotide #	NM_020630.4
NCBI Official Full Name	proto-oncogene tyrosine-protein kinase receptor Ret isoform c
NCBI Official Synonym Full Names	ret proto-oncogene
NCBI Official Symbol	RET
NCBI Official Synonym Symbols	PTC; MTC1; HSCR1; MEN2A; MEN2B; RET51; CDHF12; CDHR16; RET-ELE1
UniProt Protein Name	Proto-oncogene tyrosine-protein kinase receptor Ret
UniProt Synonym Protein Names	Cadherin family member 12; Proto-oncogene c-Ret
Protein Family	Proto-oncogene tyrosine-protein kinase receptor
UniProt Gene Name	RET
UniProt Synonym Gene Names	CDHF12; CDHR16; PTC; RET51

UniProt Entry Name	RET_HUMAN
NCBI Protein Information	proto-oncogene tyrosine-protein kinase receptor Ret; proto-oncogene c-Ret; receptor tyrosine kinase; RET transforming sequence; cadherin family member 12; hydroxyaryl-protein kinase; cadherin-related family member 16; ret proto-oncogene (multiple endocrin
NCBI Summary for RET	This gene, a member of the cadherin superfamily, encodes one of the receptor tyrosine kinases, which are cell-surface molecules that transduce signals for cell growth and differentiation. This gene plays a crucial role in neural crest development, and it can undergo oncogenic activation in vivo and in vitro by cytogenetic rearrangement. Mutations in this gene are associated with the disorders multiple endocrine neoplasia, type IIA, multiple endocrine neoplasia, type IIB, Hirschsprung disease, and medullary thyroid carcinoma. Two transcript variants encoding different isoforms have been found for this gene. Additional transcript variants have been described but their biological validity has not been confirmed. [provided by RefSeq, Jul 2008]
UniProt Comments for RET	Ret: a proto-oncogenic receptor tyrosine kinase. Receptor for glial cell line-derived neurotropic factor (GDNF) and its congeners neurturin, persephin and artemin. Part of a multicompetent receptor complex with other membrane-bound ligand-binding GDNF family receptors (aGFRs). Required for development of the kidney and neural crest-derived cell types. Three alternatively spliced isoforms have been described. Protein type: Membrane protein, integral; Oncoprotein; Protein kinase, tyrosine (receptor); EC 2.7.10.1; Protein kinase, TK; Kinase, protein; TK group; Ret family Chromosomal Location of Human Ortholog: 10q11.2 Cellular Component: intracellular membrane-bound organelle; integral to plasma membrane; cytoplasm; plasma membrane; endosome membrane; receptor complex; lipid raft Molecular Function: protein binding; protein-tyrosine kinase activity; receptor activity; transmembrane receptor protein tyrosine kinase activity; calcium ion binding; ATP binding Biological Process: response to drug; regulation of cell adhesion; caspase activation; positive regulation of neuron maturation; posterior midgut development; peptidyl-tyrosine phosphorylation; positive regulation of cell size; positive regulation of transcription, DNA-dependent; membrane protein proteolysis; MAPKKK cascade; positive regulation of cell adhesion mediated by integrin; response to pain; regulation of axonogenesis; signal transduction; neuron maturation; protein amino acid phosphorylation; enteric nervous system development; retina development in camera-type eye; ureteric bud development; embryonic epithelial tube formation; neuron adhesion; homophilic cell adhesion; neural crest cell migration; transmembrane receptor protein tyrosine kinase signaling pathway; positive regulation of cell migration Disease: Multiple Endocrine Neoplasia, Type Iib; Multiple Endocrine Neoplasia, Type Iia; Thyroid Carcinoma, Papillary; Hirschsprung Disease, Susceptibility To, 1; Thyroid Carcinoma, Familial Medullary
Research Articles on RET	**1.** The G691S variant allele does increase the risk for MTC, with a recessive mechanism of action, apparently more evident among females.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.

Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/ret/6341394