SIRT3 Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	SIRT3, Polyclonal Antibody
Full Product Name	SIRT3, NT (SIRT3, SIR2L3, NAD-dependent protein deacetylase sirtuin-3, mitochondrial, Regulatory protein SIR2 homolog 3, SIR2-like protein 3) (MaxLight 550)
Product Gene Name	anti-SIRT3 antibody
Product Synonym Names	SIRT3; SIR2L3; NAD-dependent protein deacetylase sirtuin-3, mitochondrial; Regulatory protein SIR2 homolog 3; SIR2-like protein 3
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight550.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight550 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	28,567 Da
OMIM	604481
3D Structure	ModBase 3D Structure for Q9NTG7
Immunogen	SIRT3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 55-84 amino acids from the N-terminal region of human SIRT3.
Conjugate	MaxLight550
Note	Preservative Free
Product Categories/Family for anti-SIRT3 antibody	Antibodies; Mitochondrial Proteins
Applications Tested/Suitable for anti-SIRT3 antibody	Western Blot (WB), FLISA
Application Notes for anti-SIRT3 antibody	WB: 1:100-500 FLISA: 1:1,000 Applications are based on unconjugated antibody.
Related Product Information for anti-SIRT3 antibody	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Two alternatively spliced transcript variants that encode different proteins have been described for this gene.

Product References and Citations for anti-SIRT3 antibody	Dransfeld, C.L., et al. Int. J. Oncol. 36(4):955-960(2010). Shulga, N., et al. J. Cell. Sci. 123 (PT 6), 894-902 (2010): Shi, T., et al. J Dig Dis 11(1):55-62(2010). Kong, X., et al. PLoS ONE 5 (7), E11707 (2010): Li, S., et al. PLoS ONE 5 (5), E10486 (2010).
Other Notes	Small volumes of anti-SIRT3 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for SIRT3 purchase	MBS6347809 is an antibody targeting SIRT3. Also known as SIRT3, NT (SIRT3, SIR2L3, NAD-dependent protein deacetylase sirtuin-3, mitochondrial, Regulatory protein SIR2 homolog 3, SIR2-like protein 3) (MaxLight 550). Reactive in Human. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for SIRT3. It may not necessarily be applicable to this product.

NCBI GI #	6912660
NCBI GeneID	23410
NCBI Accession #	NP_036371.1
NCBI Related Accession #	NP_001017524.1
UniProt Primary Accession #	Q9NTG7
UniProt Secondary Accession #	Q9Y6E8; B7Z5U6
UniProt Related Accession #	Q9NTG7
GenBank Nucleotide #	NM_012239.5
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-3, mitochondrial isoform a
NCBI Official Synonym Full Names	sirtuin 3
NCBI Official Symbol	SIRT3
NCBI Official Synonym Symbols	SIR2L3
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-3, mitochondrial
UniProt Synonym Protein Names	Regulatory protein SIR2 homolog 3; SIR2-like protein 3
Protein Family	NAD-dependent protein deacetylase sirtuin
UniProt Gene Name	SIRT3
UniProt Synonym Gene Names	SIR2L3; hSIRT3
UniProt Entry Name	SIR3_HUMAN
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-3, mitochondrial

UniProt Comments for SIRT3	SIRT3: a soluble mitochondrial NAD-dependent protein deacetylase. Can deacetylate and thereby activate a central metabolic regulator in the mitochondrial matrix, glutamate dehydrogenase. Mitochondrial proteins are hyperacetylated in SIRT3-deficient mice, but not in SIRT4 or SIRT5 deficient mice. SIRT3-deficient mice have no obvious metabolic phenotype, and the process of adaptive thermogenesis is normal. Can deacetylate and activate isocitrate dehydrogenase 2, an enzyme that participates in regulation of the citric acid cycle and the regeneration of antioxidants. Inhibited by nicotinamide, but not by trichostatin A. Its unprocessed form is enzymatically inactive. It is processed by mitochondrial processing peptidase (MPP) in the mitochondrial matrix to give an enzymatically active 28 kDa product. Protein type: Deacetylase; EC 3.5.1.-; Mitochondrial Chromosomal Location of Human Ortholog: 11p15.5 Cellular Component: mitochondrion; membrane; mitochondrial matrix Molecular Function: protein binding; enzyme binding; zinc ion binding; NAD-dependent histone deacetylase activity (H3-K14 specific); NAD+ ADP-ribosyltransferase activity Biological Process: mitochondrion organization and biogenesis; protein amino acid ADP-ribosylation; protein amino acid deacetylation; organelle organization and biogenesis; aerobic respiration; aging
Research Articles on SIRT3	**1.** Development of novel SIRT3 inhibitors, such as LC-0296, might enable the development of new targeted therapies to treat and improve the survival rate of patients with head and neck cancer.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/sirt3/6347809