
SRC Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	SRC, Polyclonal Antibody
Full Product Name	SRC, NT (SRC, SRC1, Proto-oncogene tyrosine-protein kinase Src, Proto-oncogene c-Src, pp60c-src) (FITC)
Product Gene Name	anti-SRC antibody
Product Synonym Names	SRC; SRC1; Proto-oncogene tyrosine-protein kinase Src; Proto-oncogene c-Src; pp60c-src
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with Fluorescein Isothiocyanate (FITC).
Purity/Purification	Purified by Protein A Affinity Chromatography.
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C. Aliquots are stable at -20 degree C for 12 months after receipt. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: FITC conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.
Molecular Weight	59,835 Da
Chromosome Location	Chromosome: 20; NC_000020.10 (35973088..36033821). Location: 20q12-q13
OMIM	190090
3D Structure	ModBase 3D Structure for P12931
Immunogen	SRC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 24-52 amino acids from the N-terminal region of human SRC.
Conjugate	FITC
Note	Preservative Free
Special Handling	Light sensitive
Product Categories/Family for anti-SRC antibody	Antibodies; Protein Kinases
Applications Tested/Suitable for anti-SRC antibody	Western Blot (WB), FLISA
Application Notes for anti-SRC antibody	WB: 1:100-500 FLISA: 1:1,000 Applications are based on unconjugated antibody.
Related Product Information for anti-SRC antibody	This gene is highly similar to the v-src gene of Rous sarcoma virus. This proto-oncogene may play a role in the regulation of embryonic development and cell growth. The protein encoded by this gene is a tyrosine-protein kinase whose activity can be inhibited by phosphorylation by c-SRC kinase. Mutations in this gene could be involved in the malignant progression of colon cancer. Two transcript variants encoding the same protein have been found for this gene.

Product References and Citations for anti-SRC antibody	Pradhan, S., et al. J. Biol. Chem. 285(38):29059-29068(2010). Dergai, M., et al. Biochem. Biophys. Res. Commun. 399(2):307-312(2010). Liu, H., et al. J. Biol. Chem. 285(29):21934-21942(2010). Pene-Dumitrescu, T., et al. J. Biol. Chem. 285(28):21446-21457(2010). Liu, X., et al. Acta Biochim. Biophys. Sin. (Shanghai) 42(7):496-501(2010).
Other Notes	Small volumes of anti-SRC antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for SRC purchase	MBS6351062 is an antibody targeting SRC. Also known as SRC, NT (SRC, SRC1, Proto-oncogene tyrosine-protein kinase Src, Proto-oncogene c-Src, pp60c-src) (FITC). Reactive in Human. Host: Rabbit.



Western Blot (WB) of anti-SRC antibody
Western blot analysis of SRC Antibody (N-term) Pab pre-incubated without(lane 1) and with(lane 2) blocking peptide in MCF-7 cell line lysate. SRC Antibody (N-term) (arrow) was detected using the purified Pab.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for SRC. It may not necessarily be applicable to this product.

NCBI GI #	4885609
NCBI GeneID	6714
NCBI Accession #	NP_005408.1
UniProt Primary Accession #	P12931
UniProt Secondary Accession #	Q76P87; Q86VB9; Q9H5A8; E1P5V4
UniProt Related Accession #	P12931
GenBank Nucleotide #	NM_005417.3
NCBI Official Full Name	proto-oncogene tyrosine-protein kinase Src
NCBI Official Synonym Full Names	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)
NCBI Official Symbol	SRC
NCBI Official Synonym Symbols	ASV; SRC1; c-SRC; p60-Src
UniProt Protein Name	Proto-oncogene tyrosine-protein kinase Src
UniProt Synonym Protein Names	Proto-oncogene c-Src; pp60c-src
Protein Family	Neuronal proto-oncogene tyrosine-protein kinase
UniProt Gene Name	SRC
UniProt Synonym Gene Names	SRC1; p60-Src
UniProt Entry Name	SRC_HUMAN

NCBI Protein Information	proto-oncogene tyrosine-protein kinase Src; proto-oncogene c-Src; tyrosine kinase pp60c-src; tyrosine-protein kinase SRC-1; protooncogene SRC, Rous sarcoma
UniProt Comments for SRC	Src: proto-oncogenic cytoplasmic tyrosine kinase of the SRC family. Highly expressed in certain fully differentiated cells such as neurons, platelets and macrophages. Phosphorylation of an activation loop tyrosine activates the enzyme; phosphorylation of a tyrosine in the C-terminus by Csk inhibits the enzyme. Two alternatively spliced isoforms have been described. Protein type: Protein kinase, TK; EC 2.7.10.2; Kinase, protein; Oncoprotein; Protein kinase, tyrosine (non-receptor); TK group; Src family Chromosomal Location of Human Ortholog: 20q12-q13 Cellular Component: neuron projection; mitochondrion; lysosome; postsynaptic density; actin filament; caveola; cytosol; extrinsic to internal side of plasma membrane; perinuclear region of cytoplasm; late endosome; mitochondrial inner membrane; cytoplasm; plasma membrane; nucleus Molecular Function: protein C-terminus binding; ephrin receptor binding; non-membrane spanning protein tyrosine kinase activity; phosphoprotein binding; insulin receptor binding; protein kinase activity; integrin binding; protein binding; enzyme binding; SH3/SH2 adaptor activity; protein kinase C binding; protein-tyrosine kinase activity; heme binding; estrogen receptor binding; SH2 domain binding; kinase activity; ATP binding; hormone receptor binding; receptor binding Biological Process: oogenesis; regulation of estrogen receptor signaling pathway; progesterone receptor signaling pathway; central nervous system development; positive regulation of cyclin-dependent protein kinase activity; estrogen receptor signaling pathway; viral reproduction; nerve growth factor receptor signaling pathway; positive regulation of apoptosis; regulation of cell cycle; positive regulation of transcription, DNA-dependent; uterus development; negative regulation of mitochondrial depolarization; negative regulation of protein homooligomerization; positive regulation of MAP kinase activity; cell-cell adhesion; transforming growth factor beta receptor signaling pathway; cell adhesion; response to electrical stimulus; bone resorption; response to drug; platelet activation; fibroblast growth factor receptor signaling pathway; activation of protein kinase B; response to virus; transcytosis; positive regulation of integrin activation; positive regulation of protein amino acid autophosphorylation; cellular response to insulin stimulus; response to mechanical stimulus; T cell costimulation; regulation of vascular permeability; negative regulation of transcription, DNA-dependent; leukocyte migration; negative regulation of apoptosis; axon guidance; peptidyl-tyrosine phosphorylation; protein amino acid autophosphorylation; platelet-derived growth factor receptor signaling pathway; negative regulation of caspase activity; signal transduction; positive regulation of smooth muscle cell migration; regulation of cell-cell adhesion; forebrain development; ephrin receptor signaling pathway; epidermal growth factor receptor signaling pathway; integrin-mediated signaling pathway; response to nutrient levels; regulation of bone resorption; negative regulation of focal adhesion formation; signal complex assembly; positive regulation of phosphoinositide 3-kinase activity; response to mineralocorticoid stimulus; cell cycle; regulation of cell proliferation; cell proliferation; positive regulation of protein kinase B signaling cascade; peptidyl-serine phosphorylation; response to hydrogen peroxide; regulation of protein binding; Ras protein signal transduction; stress fiber formation; innate immune response; response to acidity; positive regulation of insulin receptor signaling pathway; vascular endothelial growth factor receptor signaling pathway; blood coagulation; positive regulation of cytokine secretion
Research Articles on SRC	**1.** Src family kinases could play a pivotal role in increasing B cell receptor sensitivity to monomeric antigens and in mediating a rapid response to soluble multimeric antigens of pathogens that can induce spatial BCR clustering.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.

Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/src/6351062