VEGFA Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	VEGFA, Polyclonal Antibody
Full Product Name	VEGFA, ID (VEGFA, VEGF, Vascular endothelial growth factor A, Vascular permeability factor) (MaxLight 750)
Product Gene Name	anti-VEGFA antibody
Product Synonym Names	VEGFA; VEGF; Vascular endothelial growth factor A; Vascular permeability factor
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight750.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Specificity	Recognizes human VEGFA.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight750 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	27kDa
Chromosome Location	Chromosome: 6; NC_000006.11 (43737946..43754224). Location: 6p12
OMIM	192240
3D Structure	ModBase 3D Structure for P15692
Immunogen	KLH-conjugated synthetic peptide mapping to a fragment of residues within amino acids 140-168 in the Central region of human VEGFA, UniProt Accession #NP_001020537.
Conjugate	MaxLight750
Note	Preservative Free
Product Categories/Family for anti-VEGFA antibody	Antibodies; Growth Factors; Cytokines
Applications Tested/Suitable for anti-VEGFA antibody	Western Blot (WB), FLISA
Application Notes for anti-VEGFA antibody	WB: 1:1000 Applications are based on unconjugated antibody.
Related Product Information for anti-VEGFA antibody	MBS6362453 is an antibody targeting VEGFA. Also known as VEGFA, ID (VEGFA, VEGF, Vascular endothelial growth factor A, Vascular permeability factor) (MaxLight 750). Reactive in Human. Host: Rabbit.
Product References and Citations for anti-VEGFA antibody	Shrivastava-Ranjan, P., et al. J. Virol. 84(21):11227-11234(2010). Kim, Y.H., et al. Gynecol. Oncol. 119(2):232-236(2010). Yang, Y., et al. Exp. Biol. Med. (Maywood) 235(10):1204-1211(2010). Huez, I., et al. Mol. Endocrinol. 15(12):2197-2210(2001). Tee, M.K., et al. Biochem. J. 359 (PT 1), 219-226 (2001).

Other Notes	Small volumes of anti-VEGFA antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for VEGFA. It may not necessarily be applicable to this product.

NCBI GI #	76781480
NCBI GeneID	7422
NCBI Accession #	NP_001020537.2
NCBI Related Accession #	NP_001020538.2
UniProt Primary Accession #	P15692
UniProt Secondary Accession #	O60720; O75875; Q074Z4; Q16889; Q5UB46; Q6P0P5; Q96KJ0; Q96L82; Q96NW5; Q9H1W8; B5BU86
UniProt Related Accession #	P15692
GenBank Nucleotide #	NM_001025366.2
NCBI Official Full Name	vascular endothelial growth factor A isoform a
NCBI Official Synonym Full Names	vascular endothelial growth factor A
NCBI Official Symbol	VEGFA
NCBI Official Synonym Symbols	VPF; VEGF; MVCD1
UniProt Protein Name	Vascular endothelial growth factor A
UniProt Synonym Protein Names	Vascular permeability factor
UniProt Gene Name	VEGFA
UniProt Synonym Gene Names	VEGF; VEGF-A; VPF
UniProt Entry Name	VEGFA_HUMAN
NCBI Protein Information	vascular endothelial growth factor A; vascular permeability factor

UniProt Comments for VEGFA	Function: Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. Ref.27 Ref.30 Ref.33 Subunit structure: Homodimer; disulfide-linked. Also found as heterodimer with PGF By similarity. Subcellular location: Secreted. Note: VEGF121 is acidic and freely secreted. VEGF165 is more basic, has heparin-binding properties and, although a signicant proportion remains cell-associated, most is freely secreted. VEGF189 is very basic, it is cell-associated after secretion and is bound avidly by heparin and the extracellular matrix, although it may be released as a soluble form by heparin, heparinase or plasmin. Ref.26 Ref.28 Ref.32 Tissue specificity: Isoform VEGF189, isoform VEGF165 and isoform VEGF121 are widely expressed. Isoform VEGF206 and isoform VEGF145 are not widely expressed. Induction: Regulated by growth factors, cytokines, gonadotropins, nitric oxide, hypoxia, hypoglycemia and oncogenic mutations. Involvement in disease: Microvascular complications of diabetes 1 (MVCD1) [MIM:603933]: Pathological conditions that develop in numerous tissues and organs as a consequence of diabetes mellitus. They include diabetic retinopathy, diabetic nephropathy leading to end-stage renal disease, and diabetic neuropathy. Diabetic retinopathy remains the major cause of new-onset blindness among diabetic adults. It is characterized by vascular permeability and increased tissue ischemia and angiogenesis.Note: Disease susceptibility is associated with variations affecting the gene represented in this entry. Sequence similarities: Belongs to the PDGF/VEGF growth factor family. Sequence caution: The sequence AAC63102.1 differs from that shown. Reason: Erroneous initiation. The sequence AAC63143.1 differs from that shown. Reason: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon.The sequence CAC19512.2 differs from that shown. Reason: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon.The sequence CAC19516.2 differs from that shown. Reason: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/vegfa/6362453