
CDK5 Antibody – Rabbit Polyclonal


PRODUCT OVERVIEW

Product Name	CDK5, Polyclonal Antibody
Full Product Name	CDK5 (Cyclin-Dependent Kinase 5, Cell Division Protein Kinase 5, Serine/Threonine-protein Kinase PSSALRE, Tau Protein Kinase II Catalytic Subunit
Product Gene Name	anti-CDK5 antibody
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with Biotin.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Specificity	Recognizes human CDK5.
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and stor cap.
Molecular Weight	33,304 Da
Chromosome Location	Chromosome: 7; NC_000007.13 (150750899..150755052, complement). Location: 7q36
OMIM	123831
3D Structure	ModBase 3D Structure for Q00535
Immunogen	Full length human CDK5, aa1-292 (NP_004926.1).
Immunogen Sequence	MQKYEKLEKIGEGTYGTVFKAKNRETHEIVALKRVRLDDDDEGVPSSALREICLLKELKHKNIVRLHDVLHSDKKLTLVFEFCDQDLKKYFDSCNGDLDPEI
Conjugate	Biotin
Note	Preservative Free
Product Categories/Family for anti-CDK5 antibody	Antibodies; Protein Kinases
Applications Tested/Suitable for anti-CDK5 antibody	Western Blot (WB)
Application Notes for anti-CDK5 antibody	Applications are based on unconjugated antibody.

Related Product Information for anti-CDK5 antibody	Cdk5 is ubiquitously expressed, but expression and activity is highest in post mitotic neurones, where p35 and p39 are highly expressed. Deregulation
Other Notes	Small volumes of anti-CDK5 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necess
Searchable Terms for CDK5 purchase	MBS6373545 is an antibody targeting CDK5. Also known as CDK5 (Cyclin-Dependent Kinase 5, Cell Division Protein Kinase 5, Serine/Threonine-prote



Western Blot (WB) of anti-CDK5 antibody
CDK5 rabbit polyclonal antibody. Western Blot analysis of CDK5 expression in HeLa.



Western Blot (WB) of anti-CDK5 antibody
Western Blot analysis of CDK5 expression in transfected 293T cell line by CDK5 polyclonal antibody. Lane 1: CDK5 transfected lysate (33.3kD). Lane 2: Non-transfected lysate.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for CDK5. It may not necessarily be applicable to this product.

NCBI GI #	4826675
NCBI GeneID	1020
NCBI Accession #	NP_004926.1
UniProt Primary Accession #	Q00535
UniProt Secondary Accession #	A1XKG3
UniProt Related Accession #	Q00535
GenBank Nucleotide #	NM_004935.2
NCBI Official Full Name	cyclin-dependent kinase 5 isoform 1
NCBI Official Synonym Full Names	cyclin-dependent kinase 5
NCBI Official Symbol	CDK5
NCBI Official Synonym Symbols	PSSALRE
UniProt Protein Name	Cyclin-dependent kinase 5
UniProt Synonym Protein Names	Cell division protein kinase 5; Serine/threonine-protein kinase PSSALRE; Tau protein kinase II catalytic subunit
Protein Family	Cyclin-dependent kinase
UniProt Gene Name	CDK5
UniProt Synonym Gene Names	CDKN5; TPKII catalytic subunit
UniProt Entry Name	CDK5_HUMAN

NCBI Protein Information	cyclin-dependent kinase 5; TPKII catalytic subunit; protein kinase CDK5 splicing; cell division protein kinase 5; serine/threonine-protein kinase PSSALRE; tau protein kinase II catalytic subunit
NCBI Summary for CDK5	This gene encodes a proline-directed serine/threonine kinase that is a member of the cyclin-dependent kinase family of proteins. Unlike other members of the family, the protein encoded by this gene does not directly control cell cycle regulation. Instead the protein, which is predominantly expressed at high levels in mammalian postmitotic central nervous system neurons, functions in diverse processes such as synaptic plasticity and neuronal migration through phosphorylation of proteins required for cytoskeletal organization, endocytosis and exocytosis, and apoptosis. In humans, an allelic variant of the gene that results in undetectable levels of the protein has been associated with lethal autosomal recessive lissencephaly-7. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2015]
UniProt Comments for CDK5	CDK5: a protein kinase of the CDK family. Unlike other members of this family, it is not activated by cyclins but by p35 (CDK5R1) and p39. An important regulator of neuronal positioning during brain development. May also play a role in synaptogenesis and neurotransmission. Substrates include TAU, MAP2, NF-H and -M, Nudel, PDE6, beta-catenin, amphyphysin, dynamin I, synapsin 1, Munc-18, and NMDA receptor 2A. Plays a role in myogenesis, haematopoietic cell differentiation, spermatogenesis, insulin secretion, and lens differentiation. Implicated in the pathology of neurofibrillary tangles and formation of senile plaques, hallmarks of Alzheimer?s disease. Induces tau phosphorylation and aggregation and neurofibrillary tangle deposition and neurodegeneration in in vitro and in vivo animal models. Brain samples from Alzeimer?s pateints show elevated CDK5 activity. Protein type: Kinase, protein; EC 2.7.11.1; Cell cycle regulation; Protein kinase, Ser/Thr (non-receptor); Protein kinase, CMGC; CMGC group; CDK family; CDK5 subfamily; CDK/CDK5 subfamily Chromosomal Location of Human Ortholog: 7q36 Cellular Component: cyclin-dependent protein kinase 5 activator complex; dendrite; postsynaptic density; perikaryon; cytosol; postsynaptic membrane; cytoskeleton; growth cone; membrane; cell soma; axon; lamellipodium; cytoplasm; nucleus; cell junction; neuromuscular junction; filopodium Molecular Function: acetylcholine receptor activator activity; ephrin receptor binding; p53 binding; protein kinase activity; protein serine/threonine kinase activity; ErbB-2 class receptor binding; protein binding; ErbB-3 class receptor binding; cyclin-dependent protein kinase activity; cytoskeletal protein binding; tau-protein kinase activity; kinase activity; ATP binding Biological Process: Schwann cell development; negative regulation of synaptic plasticity; axon extension; cerebellar cortex formation; rhythmic process; regulation of synaptic plasticity; motor axon guidance; sensory perception of pain; receptor clustering; regulation of cell migration; corpus callosum development; regulation of apoptosis; neuron differentiation; receptor catabolic process; oligodendrocyte differentiation; neurite development; central nervous system neuron development; synaptic transmission, glutamatergic; skeletal muscle development; dendrite morphogenesis; cell division; synaptic vesicle endocytosis; negative regulation of protein ubiquitination; negative regulation of transcription, DNA-dependent; axon guidance; regulated secretory pathway; synaptic transmission, dopaminergic; negative regulation of protein export from nucleus; positive regulation of protein binding; protein amino acid autophosphorylation; cell-matrix adhesion; neuron migration; nucleocytoplasmic transport; negative regulation of axon extension; negative regulation of cell cycle; synaptic transmission; intracellular protein transport; synaptogenesis; behavioral response to cocaine; positive regulation of neuron apoptosis; visual learning; cortical actin cytoskeleton organization and biogenesis; layer formation in the cerebral cortex; serine phosphorylation of STAT3 protein; negative regulation of proteolysis; hippocampus development; peptidyl-threonine phosphorylation; cell cycle; cell proliferation; positive regulation of calcium ion-dependent exocytosis; embryonic development; synaptic vesicle exocytosis; peptidyl-serine phosphorylation; neuron apoptosis; positive regulation of protein kinase activity; blood coagulation; regulation of excitatory postsynaptic membrane potential; phosphorylation Disease: Lissencephaly 7 With Cerebellar Hypoplasia
Research Articles on CDK5	**1.** UCH-L1 physically interacts with CDK1, CDK4, and CDK5, enhancing their kinase activity.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.

Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/cdk5/6373545