MEF2C Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	MEF2C, Polyclonal Antibody
Full Product Name	MEF2C (S59) (Myocyte-specific Enhancer Factor 2C) (MaxLight 490)
Product Gene Name	anti-MEF2C antibody
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight490.
Purity/Purification	Purified by Immunoaffinity chromatography.
Specificity	Recognizes human MEF2C.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight490 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	50,242 Da
OMIM	600662
3D Structure	ModBase 3D Structure for Q06413
Immunogen	Synthetic phosphopeptide corresponding to amino acid residues surrounding Thr300 of human MEF2C (KLH).
Conjugate	MaxLight490
Note	Preservative Free
Product Categories/Family for anti-MEF2C antibody	Antibodies; Transcription Factors
Applications Tested/Suitable for anti-MEF2C antibody	Western Blot (WB), FLISA
Application Notes for anti-MEF2C antibody	WB: 1:50-1:100 FLISA: 1:1,000 Applications are based on unconjugated antibody.
Related Product Information for anti-MEF2C antibody	MBS6488989 is an antibody targeting MEF2C. Also known as MEF2C (S59) (Myocyte-specific Enhancer Factor 2C) (MaxLight 490). Reactive in Human. Host: Rabbit.
Product References and Citations for anti-MEF2C antibody	1. Konig, S., et al., J. Biol. Chem. 279(27):28187-28196 (2004). 2. Maeda, T., et al., J. Biol. Chem. 277(50):48889-48898 (2002). 3. Maeda, T., et al., Biochem. Biophys. Res. Commun. 294(4):791-797 (2002). 4. Janson, C.G., et al., Brain Res. Mol. Brain Res. 97(1):70-82 (2001). 5. Krainc, D., et al., Genomics 29(3):809-811 (1995).
Other Notes	Small volumes of anti-MEF2C antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for MEF2C. It may not necessarily be applicable to this product.

NCBI GI #	19923215
NCBI GeneID	4208
NCBI Accession #	NP_002388
UniProt Primary Accession #	Q06413
UniProt Secondary Accession #	C9JMZ0; D7F7N5; F8W7V7
UniProt Related Accession #	Q06413
GenBank Nucleotide #	NM_002397
NCBI Official Full Name	myocyte-specific enhancer factor 2C isoform 1
NCBI Official Synonym Full Names	myocyte enhancer factor 2C
NCBI Official Symbol	MEF2C
NCBI Official Synonym Symbols	DEL5q14.3; C5DELq14.3
UniProt Protein Name	Myocyte-specific enhancer factor 2C
Protein Family	Myocyte-specific enhancer factor
UniProt Gene Name	MEF2C
UniProt Entry Name	MEF2C_HUMAN
NCBI Protein Information	myocyte-specific enhancer factor 2C; MADS box transcription enhancer factor 2, polypeptide C

UniProt Comments for MEF2C	MEF2C: transcription factor of the MADS family which binds specifically to the MEF2 element present in the regulatory regions of many muscle-specific genes. May be involved in myogenesis, neurogenesis and in the development of cortical architecture. Three splice-variant isoforms have been described. Protein type: DNA-binding; Transcription factor Chromosomal Location of Human Ortholog: 5q14.3 Cellular Component: nucleoplasm; protein complex; intracellular membrane-bound organelle; cytoplasm; nuclear speck; nucleus; cytosol Molecular Function: protein binding; miRNA binding; transcription activator binding; DNA binding; AT DNA binding; protein heterodimerization activity; histone deacetylase binding; chromatin binding; transcription factor activity Biological Process: cardiac muscle hypertrophy; muscle development; nerve growth factor receptor signaling pathway; positive regulation of transcription, DNA-dependent; heart development; regulation of neurotransmitter secretion; stress-activated MAPK cascade; regulation of synaptic plasticity; toll-like receptor 3 signaling pathway; toll-like receptor 5 signaling pathway; neuron differentiation; positive regulation of MAP kinase activity; B cell receptor signaling pathway; regulation of neuron apoptosis; negative regulation of ossification; positive regulation of B cell proliferation; chondrocyte differentiation; negative regulation of neuron apoptosis; toll-like receptor 4 signaling pathway; positive regulation of cardiac muscle cell proliferation; endochondral ossification; nervous system development; skeletal muscle development; monocyte differentiation; response to virus; myotube differentiation; positive regulation of skeletal muscle development; muscle cell fate determination; toll-like receptor 2 signaling pathway; muscle cell differentiation; regulation of synaptogenesis; neuron development; positive regulation of transcription from RNA polymerase II promoter; smooth muscle cell differentiation; toll-like receptor 9 signaling pathway; B cell proliferation; transcription from RNA polymerase II promoter; regulation of synaptic transmission, glutamatergic; apoptosis; neuron migration; ventricular cardiac muscle cell differentiation; regulation of synaptic activity; negative regulation of transcription from RNA polymerase II promoter; palate development; toll-like receptor 10 signaling pathway; learning and/or memory; B cell homeostasis; regulation of transcription, DNA-dependent; melanocyte differentiation; regulation of megakaryocyte differentiation; germinal center formation; neural crest cell differentiation; heart looping; negative regulation of epithelial cell proliferation; blood vessel development; MyD88-independent toll-like receptor signaling pathway; MAPKKK cascade; embryonic viscerocranium morphogenesis; positive regulation of bone mineralization; humoral immune response; regulation of germinal center formation; positive regulation of myoblast differentiation; osteoblast differentiation; MyD88-dependent toll-like receptor signaling pathway; positive regulation of osteoblast differentiation; neuron morphogenesis during differentiation; toll-like receptor signaling pathway; innate immune response; positive regulation of muscle cell differentiation; blood vessel remodeling; platelet formation; regulation of excitatory postsynaptic membrane potential; positive regulation of neuron differentiation Disease: Mental Retardation, Autosomal Dominant 20
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/mef2c/6488989