
PLA2G4A Antibody – Rabbit Polyclonal


PRODUCT OVERVIEW

Product Name	PLA2G4A, Polyclonal Antibody
Full Product Name	PLA2G4A, ID (Cytosolic Phospholipase A2, cPLA2, Phospholipase A2 Group IVA, Phospholipase A2, Phosphatidylcholine 2-acylhydrolase, Lysophospholipase, PLA2G4, CPLA2) (MaxLight 490)
Product Gene Name	anti-PLA2G4A antibody
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight490.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Specificity	Recognizes human PLA2G4A.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight490 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	110kDa
Chromosome Location	Chromosome: 1; NC_000001.10 (186798032..186958113). Location: 1q25
OMIM	600522
3D Structure	ModBase 3D Structure for P47712
Immunogen	KLH-conjugated synthetic peptide mapping to a fragment of residues within amino acids 513-541 in the central region of human PLA2G4A, UniProt Accession #NP_077734.1; P47712. Species sequence homology: Chicken, mouse, rat, zebrafish
Conjugate	MaxLight490
Note	Preservative Free
Product Categories/Family for anti-PLA2G4A antibody	Antibodies; Enzymes; Phospholipase
Applications Tested/Suitable for anti-PLA2G4A antibody	Western Blot (WB), FLISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Flow Cytometry (FC/FACS)
Application Notes for anti-PLA2G4A antibody	WB: 1:100-1:500 FLISA: 1:1,000 Applications are based on unconjugated antibody.
Related Product Information for anti-PLA2G4A antibody	MBS6493271 is an antibody targeting PLA2G4A. Also known as PLA2G4A, ID (Cytosolic Phospholipase A2, cPLA2, Phospholipase A2 Group IVA, Phospholipase A2, Phosphatidylcholine 2-acylhydrolase, Lysophospholipase, PLA2G4, CPLA2) (MaxLight 490). Reactive in Human. Host: Rabbit.
Product References and Citations for anti-PLA2G4A antibody	1. Sharp, J.D., et.al., J. Biol. Chem. 266(23), 14850-14853 (1991). 2. Clark, J.D., et.al., Cell 65(6), 1043-1051 (1991).


Other Notes	Small volumes of anti-PLA2G4A antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.




Western Blot (WB) of anti-PLA2G4A antibody
Western Blot analysis in NCI-H460 cell line lysates (35ug/lane) usingMBS625712. PLA2G4A (arrow) was detected using the purified Pab.



Immunohistochemistry (IHC) of anti-PLA2G4A antibody
Immunohistochemistry analysis in formalin-fixed and paraffin-embedded human lung carcinoma withMBS625712, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of MBS625712 for immunohistochemistry.



Immunofluorescence (IF) of anti-PLA2G4A antibody
Confocal immunofluorescent analysis in NCI-H460 cells using MBS625712 followed by Alexa Fluoro 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Flow Cytometry (FC/FACS) of anti-PLA2G4A antibody
Flow Cytometric analysis of NCI-H292 cells using MBS625712 (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for PLA2G4A. It may not necessarily be applicable to this product.

NCBI GI #	23943920
NCBI GeneID	5321
NCBI Accession #	NP_077734.1
UniProt Primary Accession #	P47712
UniProt Secondary Accession #	Q29R80; B1AKG4
UniProt Related Accession #	P47712
GenBank Nucleotide #	NM_024420.2
NCBI Official Full Name	cytosolic phospholipase A2
NCBI Official Synonym Full Names	phospholipase A2, group IVA (cytosolic, calcium-dependent)
NCBI Official Symbol	PLA2G4A

NCBI Official Synonym Symbols	PLA2G4; cPLA2-alpha
UniProt Protein Name	Cytosolic phospholipase A2
UniProt Synonym Protein Names	Phospholipase A2 group IVAIncluding the following 2 domains:Phospholipase A2 (EC:3.1.1.4)Alternative name(s):Phosphatidylcholine 2-acylhydrolase
Protein Family	Cytosolic phospholipase
UniProt Gene Name	PLA2G4A
UniProt Synonym Gene Names	CPLA2; PLA2G4; cPLA2
UniProt Entry Name	PA24A_HUMAN
NCBI Protein Information	cytosolic phospholipase A2; cPLA2; lysophospholipase; phospholipase A2 group IVA; phosphatidylcholine 2-acylhydrolase; calcium-dependent phospholipid-binding protein
NCBI Summary for PLA2G4A	This gene encodes a member of the cytosolic phospholipase A2 group IV family. The enzyme catalyzes the hydrolysis of membrane phospholipids to release arachidonic acid which is subsequently metabolized into eicosanoids. Eicosanoids, including prostaglandins and leukotrienes, are lipid-based cellular hormones that regulate hemodynamics, inflammatory responses, and other intracellular pathways. The hydrolysis reaction also produces lysophospholipids that are converted into platelet-activating factor. The enzyme is activated by increased intracellular Ca(2+) levels and phosphorylation, resulting in its translocation from the cytosol and nucleus to perinuclear membrane vesicles. [provided by RefSeq, Jul 2008]
UniProt Comments for PLA2G4A	cPLA2: a calcium-dependent phospholipase A2 that catalyzes the release of arachidonic acid from membrane phospholipids. Selectively hydrolyzes arachidonyl phospholipids in the sn-2 position releasing arachidonic acid. Arachidonic acid is a precursor for the eicosanoids that are involved in hemodynamic regulation, inflammatory responses, and other cellular processes. Promotes cerebellar long-term depression and motor learning. Translocates to the Golgi and endoplasmic reticulum in a calcium-dependent fashion. Translocation and activation of at the ER facilitates the process of ER stress. Regulates the biogenesis of lipid droplets, a process dependent on JNK and ceramide kinase. Stimulated by agonists such as ATP, EGF, thrombin and bradykinin as well as by cytosolic Ca2+. The N-terminal C2 domain, by its association with lipid membranes, mediates the regulation of CPLA2 by presenting the active site to its substrate in response to elevations of cytosolic Ca2+. Inhibited when in trimolecular complex with ANXA2 and S100A10. Phosphorylation of S727 relieves this inhibitory interaction, thus activating PLA2G4A. Protein type: Lipid Metabolism - ether lipid; EC 3.1.1.4; Lipid Metabolism - glycerophospholipid; Lipid Metabolism - arachidonic acid; EC 3.1.1.5; Phospholipase; Lipid Metabolism - linoleic acid; Lipid Metabolism - alpha-linolenic acid Chromosomal Location of Human Ortholog: 1q25 Cellular Component: Golgi apparatus; endoplasmic reticulum membrane; cytoplasm; cytoplasmic membrane-bound vesicle; mitochondrial inner membrane; cytosol Molecular Function: phospholipase A2 activity; calcium-dependent phospholipase A2 activity; calcium-dependent phospholipid binding; calcium ion binding; lysophospholipase activity Biological Process: icosanoid biosynthetic process; platelet activating factor biosynthetic process; platelet activation; phospholipid catabolic process; phospholipid metabolic process; glycerophospholipid biosynthetic process; icosanoid metabolic process; phosphatidic acid biosynthetic process; arachidonic acid metabolic process; arachidonic acid secretion; blood coagulation; regulation of cell proliferation
Research Articles on PLA2G4A	**1.** results suggest that C2-di-ethyl-ceramide-1-phosphate appeared to inhibit cPLA(2)alpha, probably by interaction with a site in the catalytic domain of the enzyme.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.

Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/pla2g4a/6493271