# PRKCA Antibody – Rabbit Polyclonal

## PRODUCT OVERVIEW

| | |
|---|---|
| **Product Name** | PRKCA, Polyclonal Antibody |
| **Full Product Name** | PRKCA, CT (Protein Kinase C alpha Type, AAG6, PKC-A, PRKACA, PKCA, PKC-alpha) (MaxLight 405) |
| **Product Gene Name** | anti-PRKCA antibody |
| **Clonality** | Polyclonal |
| **Isotype** | IgG |
| **Host** | Rabbit |
| **Species Reactivity** | Human |
| **Form/Format** | Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight405. |
| **Purity/Purification** | Purified by Protein G Affinity Chromatography. |
| **Specificity** | Recognizes human PKC. |
| **Preparation and Storage** | Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight405 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap. |
| **Molecular Weight** | 76,750 Da |
| **Chromosome Location** | Chromosome: 17; NC_000017.10 (64298926..64806862). Location: 17q22-q23.2 |
| **OMIM** | 176960 |
| **3D Structure** | ModBase 3D Structure for P17252 |
| **Immunogen** | Synthetic peptide selected from the C-terminal region of human PKC (KLH). |
| **Conjugate** | MaxLight405 |
| **Note** | Preservative Free |
| **Product Categories/Family for anti-PRKCA antibody** | Antibodies; Protein Kinases |
| **Applications Tested/Suitable for anti-PRKCA antibody** | Immunohistochemistry (IHC), FLISA, Western Blot (WB) |
| **Application Notes for anti-PRKCA antibody** | IHC: 1:50-1:100<br>FLISA: 1:1,000<br>WB: 1:100-1:500<br>Applications are based on unconjugated antibody. |
| **Related Product Information for anti-PRKCA antibody** | MBS6495108 is an antibody targeting PRKCA. Also known as PRKCA, CT (Protein Kinase C alpha Type, AAG6, PKC-A, PRKACA, PKCA, PKC-alpha) (MaxLight 405). Reactive in Human. Host: Rabbit. |
| **Product References and Citations for anti-PRKCA antibody** | 1. Rabinovitz, I., et al., Mol. Cell. Biol. 24(10):4351-4360 (2004). 2. Gong, N., et al., Proc. Natl. Acad. Sci. USA 101(17):6490-6495 (2004). 3. Hermoso, M., et al., J. Biol. Chem. 279(17):17681-17689 (2004). 4. Zemskov, E.A., et al., Neuroreport 14(11):1435-1438 (2003). 5. Lorenz, K., et al., Nature 426(6966):574-579 (2003). |
| **Other Notes** | Small volumes of anti-PRKCA antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply. |

## ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for PRKCA. It may not necessarily be applicable to this product.

| | |
|---|---|
| **NCBI GI #** | 4506067 |
| **NCBI GeneID** | 5578 |
| **NCBI Accession #** | NP_002728.1 |
| **UniProt Primary Accession #** | P17252 |
| **UniProt Secondary Accession #** | Q15137; Q32M72; Q96RE4; B5BU22 |
| **UniProt Related Accession #** | P17252 |
| **GenBank Nucleotide #** | NM_002737.2 |
| **NCBI Official Full Name** | protein kinase C alpha type |
| **NCBI Official Synonym Full Names** | protein kinase C, alpha |
| **NCBI Official Symbol** | PRKCA |
| **NCBI Official Synonym Symbols** | AAG6; PKCA; PRKACA; PKC-alpha |
| **UniProt Protein Name** | Protein kinase C alpha type |
| **Protein Family** | Protein kinase |
| **UniProt Gene Name** | PRKCA |
| **UniProt Synonym Gene Names** | PKCA; PRKACA; PKC-A; PKC-alpha |
| **UniProt Entry Name** | KPCA_HUMAN |
| **NCBI Protein Information** | protein kinase C alpha type; PKC-A; aging-associated gene 6 |
| **NCBI Summary for PRKCA** | Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq, Jul 2008] |

| | |
|---|---|
| **UniProt Comments for PRKCA** | PKCA: an AGC kinase of the PKC family. A classical PKC downstream of many mitogenic and receptors. Classical PKCs are calcium-dependent enzymes that are activated by phosphatidylserine, diacylglycerol and phorbol esters. Contains a pseudo-substrate autoinhibitory domain that binds to the catalytic domain preventing its activation in the absence of cofactors or activators.<br><br>Protein type: Oncoprotein; Protein kinase, AGC; Kinase, protein; EC 2.7.11.13; Protein kinase, Ser/Thr (non-receptor); AGC group; PKC family; Alpha subfamily<br><br>Chromosomal Location of Human Ortholog: 17q22-q23.2<br><br>Cellular Component: nucleoplasm; photoreceptor outer segment; cell soma; mitochondrion; perinuclear region of cytoplasm; endoplasmic reticulum; apical part of cell; cytoplasm; mitochondrial membrane; dendrite; plasma membrane; cytosol<br><br>Molecular Function: protein serine/threonine kinase activity; protein binding; enzyme binding; protein kinase C activity; zinc ion binding; calcium-dependent protein kinase C activity; ATP binding; protein kinase activity<br><br>Biological Process: phototransduction, visible light; extracellular matrix organization and biogenesis; positive regulation of cell adhesion; nerve growth factor receptor signaling pathway; regulation of muscle contraction; positive regulation of lipopolysaccharide-mediated signaling pathway; mitotic nuclear envelope disassembly; negative regulation of insulin receptor signaling pathway; signal transduction; positive regulation of mitotic cell cycle; protein amino acid phosphorylation; induction of positive chemotaxis; negative regulation of cell proliferation; synaptic transmission; chondrocyte differentiation; angiogenesis; positive regulation of macrophage differentiation; cell adhesion; regulation of peptidyl-tyrosine phosphorylation; regulation of the force of heart contraction; inactivation of MAPK activity; epidermal growth factor receptor signaling pathway; neutrophil chemotaxis; platelet activation; fibroblast growth factor receptor signaling pathway; positive regulation of blood vessel endothelial cell migration; adenylate cyclase activation; negative regulation of adenylate cyclase activity; cellular calcium ion homeostasis; rhodopsin mediated signaling; positive regulation of angiogenesis; negative regulation of glucose import; regulation of rhodopsin mediated signaling; phospholipase C activation; energy reserve metabolic process; innate immune response; positive regulation of endothelial cell proliferation; gene expression; mitotic cell cycle; positive regulation of protein amino acid phosphorylation; blood coagulation; vascular endothelial growth factor receptor signaling pathway; regulation of insulin secretion; positive regulation of inflammatory response; positive regulation of cell migration |
| **Research Articles on PRKCA** | **1.** VEGF and PKC promote degradation-independent protein ubiquitination of FLNB to control intracellular trafficking of HDAC7. |
| **Precautions** | All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines. |
| **Disclaimer** | While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice.<br><br>It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information. |
| **Product Note** | Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details. |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures.

**MyBioSource** | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/prkca/6495108