
# RYK Antibody – Rabbit Polyclonal

## PRODUCT OVERVIEW

| | |
|---|---|
| **Product Name** | RYK, Polyclonal Antibody |
| **Full Product Name** | RYK, ID (C397) (Tyrosine-protein Kinase RYK, JTK5A) (HRP) |
| **Product Gene Name** | anti-RYK antibody |
| **Clonality** | Polyclonal |
| **Isotype** | IgG |
| **Host** | Rabbit |
| **Species Reactivity** | Human |
| **Form/Format** | Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with Horseradish Peroxidase (HRP). |
| **Purity/Purification** | Purified by Ammonium Sulfate Precipitation. |
| **Specificity** | Recognizes human RYK. |
| **Preparation and Storage** | Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C. Aliquots are stable at -20 degree C for 12 months after receipt. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Note: Sodium azide is a potent inhibitor of peroxidase and should not be added to HRP conjugates. |
| **Molecular Weight** | ~66 kDa |
| **Chromosome Location** | Chromosome: 3; NC_000003.11 (133875978..133969586, complement). Location: 3q22 |
| **OMIM** | 600524 |
| **3D Structure** | ModBase 3D Structure for P34925 |
| **Immunogen** | Synthetic peptide selected from the center region of human RYK (KLH). |
| **Conjugate** | HRP |
| **Note** | Preservative Free |
| **Product Categories/Family for anti-RYK antibody** | Antibodies; Tyrosine Kinases |
| **Applications Tested/Suitable for anti-RYK antibody** | Western Blot (WB), ELISA (EIA) |
| **Application Notes for anti-RYK antibody** | WB: 1:50-1:100<br>ELISA: 1:1,000<br>Applications are based on unconjugated antibody. |
| **Related Product Information for anti-RYK antibody** | MBS6498362 is an antibody targeting RYK. Also known as RYK, ID (C397) (Tyrosine-protein Kinase RYK, JTK5A) (HRP). Reactive in Human. Host: Rabbit. |
| **Product References and Citations for anti-RYK antibody** | 1. Trivier, E., et al., J. Biol. Chem. 277(25):23037-23043 (2002). 2. Katso, R.M., et al., Mol. Cell. Biol. 19(9):6427-6440 (1999). 3. Wang, X.C., et al., Mol. Med. 2(2):189-203 (1996). 4. Tamagnone, L., et al., Oncogene 8(7):2009-2014 (1993). 5. Stacker, S.A., et al., Oncogene 8(5):1347-1356 (1993). |
| **Other Notes** | Small volumes of anti-RYK antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply. |




**Western Blot (WB) of anti-RYK antibody**
Western blot analysis of RYK antibody (Center C397) in Hela cell line lysates (35ug/lane).
RYK (arrow) was detected using the purified Pab.

## ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for RYK. It may not necessarily be applicable to this product.

| | |
|---|---|
| **NCBI GI #** | 116805325 |
| **NCBI GeneID** | 6259 |
| **NCBI Accession #** | NP_001005861.1 |
| **UniProt Primary Accession #** | P34925 |
| **UniProt Secondary Accession #** | Q04696 |
| **UniProt Related Accession #** | P34925 |
| **GenBank Nucleotide #** | NM_001005861 |
| **NCBI Official Full Name** | Homo sapiens receptor-like tyrosine kinase (RYK), transcript variant 1, mRNA |
| **NCBI Official Synonym Full Names** | receptor-like tyrosine kinase |
| **NCBI Official Symbol** | RYK |
| **NCBI Official Synonym Symbols** | JTK5; RYK1; JTK5A; D3S3195 |
| **UniProt Protein Name** | Tyrosine-protein kinase RYK |
| **Protein Family** | Tyrosine-protein kinase |
| **UniProt Gene Name** | RYK |
| **UniProt Synonym Gene Names** | JTK5A |
| **UniProt Entry Name** | RYK_HUMAN |
| **NCBI Protein Information** | tyrosine-protein kinase RYK; hydroxyaryl-protein kinase; JTK5A protein tyrosine kinase |
| **NCBI Summary for RYK** | The protein encoded by this gene is an atypical member of the family of growth factor receptor protein tyrosine kinases, differing from other members at a number of conserved residues in the activation and nucleotide binding domains. This gene product belongs to a subfamily whose members do not appear to be regulated by phosphorylation in the activation segment. It has been suggested that mediation of biological activity by recruitment of a signaling-competent auxiliary protein may occur through an as yet uncharacterized mechanism. The encoded protein has a leucine-rich extracellular domain with a WIF-type Wnt binding region, a single transmembrane domain, and an intracellular tyrosine kinase domain. This protein is involved in stimulating Wnt signaling pathways such as the regulation of axon pathfinding. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Feb 2012] |

| | |
|---|---|
| **UniProt Comments for RYK** | Function: May be a coreceptor along with FZD8 of Wnt proteins, such as WNT1, WNT3, WNT3A and WNT5A. Involved in neuron differentiation, axon guidance, corpus callosum establishment and neurite outgrowth. In response to WNT3 stimulation, receptor C-terminal cleavage occurs in its transmembrane region and allows the C-terminal intracellular product to translocate from the cytoplasm to the nucleus where it plays a crucial role in neuronal development. Ref.5<br><br>Catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.<br><br>Subunit structure: Interacts with DVL1 (via PDZ domain). Ref.5<br><br>Subcellular location: Membrane; Single-pass type I membrane protein<br><br>By similarity. Nucleus<br><br>By similarity. Cytoplasm<br><br>By similarity. Note: In cells that have undergone neuronal differentiation, the C-terminal cleaved part is translocated from the cytoplasm to the nucleus<br><br>By similarity.<br><br>Tissue specificity: Observed in all the tissues examined.<br><br>Domain: The extracellular WIF domain is responsible for Wnt binding<br><br>By similarity.<br><br>Post-translational modification: Proteolytically cleaved, in part by presenilin, in response to WNT3 stimulation. Cleavage occurs during neuronal differentiation. Ref.6<br><br>Sequence similarities: Belongs to the protein kinase superfamily. Tyr protein kinase family.Contains 1 protein kinase domain.Contains 1 WIF domain.<br><br>Caution: According to some authors, has impaired kinase activity (Ref.4). |
| **Research Articles on RYK** | **1.** RYK and VANGL2 proteins form a complex, whereas RYK also activated RhoA, a downstream effector of PCP signaling. |
| **Precautions** | All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines. |
| **Disclaimer** | While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice.<br><br>It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information. |
| **Product Note** | Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details. |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures.

**MyBioSource** | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/ryk/6498362