SMAD2 Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	Smad2, Polyclonal Antibody
Full Product Name	Smad2 (Mothers Against Decapentaplegic Homolog 2, Mothers Against DPP Homolog 2, MAD Homolog 2, Mad-related Protein 2, hMAD-2, SMAD Family Member 2, SMAD 2, hSMAD2, JV18-1, MADR2, MADH2) (MaxLight 650)
Product Gene Name	anti-Smad2 antibody
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with MaxLight650.
Purity/Purification	Purified by Immunoaffinity chromatography.
Specificity	Recognizes human SMAD2.
Preparation and Storage	Store product at 4 degree C in the dark. DO NOT FREEZE! Stable at 4 degree C for 12 months after receipt as an undiluted liquid. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: MaxLight650 conjugates are sensitive to light. For maximum recovery of product, centrifuge the original vial prior to removing the cap.
Molecular Weight	54.4 kDa (487aa)
OMIM	601366
3D Structure	ModBase 3D Structure for Q15796
Immunogen	Synthetic non-phosphopeptide corresponding to amino acid residues surrounding T220 of human SMAD2 (KLH).
Conjugate	MaxLight650
Note	Preservative Free
Product Categories/Family for anti-Smad2 antibody	Antibodies; Smad Proteins
Applications Tested/Suitable for anti-Smad2 antibody	Western Blot (WB), FLISA
Application Notes for anti-Smad2 antibody	WB: 1:50-1:100 FLISA: 1:1,000 Applications are based on unconjugated antibody.
Related Product Information for anti-Smad2 antibody	MBS6499751 is an antibody targeting Smad2. Also known as Smad2 (Mothers Against Decapentaplegic Homolog 2, Mothers Against DPP Homolog 2, MAD Homolog 2, Mad-related Protein 2, hMAD-2, SMAD Family Member 2, SMAD 2, hSMAD2, JV18-1, MADR2, MADH2) (MaxLight 650). Reactive in Human. Host: Rabbit.
Product References and Citations for anti-Smad2 antibody	1. Funaba, M., J. Biol. Chem. 277(44), 41361-41368 (2002). 2. Wicks, S.J., Mol. Cell. Biol. 20(21), 8103-8111 (2000).
Other Notes	Small volumes of anti-Smad2 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for Smad2. It may not necessarily be applicable to this product.

NCBI GI #	5174511
NCBI GeneID	4087
NCBI Accession #	NP_005892
UniProt Primary Accession #	Q15796
UniProt Related Accession #	Q15796
GenBank Nucleotide #	NM_005901.5
NCBI Official Full Name	mothers against decapentaplegic homolog 2 isoform 1
NCBI Official Synonym Full Names	SMAD family member 2
NCBI Official Symbol	SMAD2
NCBI Official Synonym Symbols	JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2
UniProt Protein Name	Mothers against decapentaplegic homolog 2
UniProt Synonym Protein Names	JV18-1; Mad-related protein 2; hMAD-2; SMAD family member 2
Protein Family	Mothers against decapentaplegic
UniProt Gene Name	SMAD2
UniProt Synonym Gene Names	MADH2; MADR2; MAD homolog 2; Mothers against DPP homolog 2; hMAD-2; SMAD 2; Smad2; hSMAD2
UniProt Entry Name	SMAD2_HUMAN
NCBI Protein Information	mothers against decapentaplegic homolog 2
NCBI Summary for Smad2	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]

UniProt Comments for Smad2	SMAD2: ubiquitously expressed transcription factor phosphorylated and activated by TGF-beta receptor-type kinases. Participates in a wide range of critical processes including morphogenesis, cell-fate determination, proliferation, differentiation and apoptosis. Phosphorylated Smads dimerize with collaborating Smad4 and are translocated into the nucleus, where the transcription of target genes is stimulated. Two alternatively spliced isoforms have been described. Protein type: DNA-binding; Transcription factor Chromosomal Location of Human Ortholog: 18q21.1 Cellular Component: nucleoplasm; transcription factor complex; cytoplasm; nuclear chromatin; nucleus; cytosol Molecular Function: transcription activator binding; metal ion binding; transcription factor binding; transforming growth factor beta receptor, pathway-specific cytoplasmic mediator activity; phosphatase binding; protein binding; DNA binding; ubiquitin protein ligase binding; double-stranded DNA binding; chromatin binding; SMAD binding; transcription factor activity; transforming growth factor beta receptor binding Biological Process: developmental growth; somatic stem cell maintenance; positive regulation of transcription, DNA-dependent; activin receptor signaling pathway; paraxial mesoderm morphogenesis; gastrulation; primary microRNA processing; negative regulation of transcription from RNA polymerase II promoter; palate development; post-embryonic development; anterior/posterior pattern formation; negative regulation of cell proliferation; regulation of transforming growth factor beta receptor signaling pathway; ureteric bud development; transforming growth factor beta receptor signaling pathway; pancreas development; embryonic foregut morphogenesis; response to glucose stimulus; positive regulation of BMP signaling pathway; pericardium development; transcription initiation from RNA polymerase II promoter; cell fate commitment; regulation of binding; transcription, DNA-dependent; in utero embryonic development; common-partner SMAD protein phosphorylation; embryonic cranial skeleton morphogenesis; SMAD protein complex assembly; organ growth; zygotic determination of dorsal/ventral axis; mesoderm formation; endoderm formation; insulin secretion; gene expression; positive regulation of transcription from RNA polymerase II promoter; negative regulation of transcription, DNA-dependent; negative regulation of transforming growth factor beta receptor signaling pathway; lung development
Research Articles on Smad2	**1.** a novel heterozygous missense mutation (c.833C>T, p.A278V) in the SMAD2 gene in a family with early onset aortic aneurysms
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/smad2/6499751