EIF3C Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	EIF3C, Polyclonal Antibody
Full Product Name	EIF3C Antibody; Biotin conjugated
Product Gene Name	anti-EIF3C antibody
Product Synonym Names	Eukaryotic translation initiation factor 3 subunit C; Eukaryotic translation initiation factor 3 subunit 8; EIF3C
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Liquid
Purity/Purification	>95%,Protein G purified
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.
Matching Pairs	Unconjugated Antibody: EIF3C (MBS969908) Biotin Conjugated Antibody: EIF3C (MBS7002795)
Molecular Weight	104,101 Da
OMIM	603916
3D Structure	ModBase 3D Structure for Q99613
Immunogen	Recombinant human Eukaryotic translation initiation factor 3 subunit C protein
Conjugation	Biotin
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-16376 / sc-31872 / sc-31873 / sc-28858
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-EIF3C antibody	ELISA (EIA)
Related Product Information for anti-EIF3C antibody	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.
Other Notes	Small volumes of anti-EIF3C antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for EIF3C purchase	MBS7002795 is an antibody targeting EIF3C. Also known as EIF3C Antibody; Biotin conjugated. Reactive in Human. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for EIF3C. It may not necessarily be applicable to this product.

NCBI GI #	83700233
NCBI GeneID	8663
NCBI Accession #	NP_001032897.1
UniProt Primary Accession #	Q99613
UniProt Secondary Accession #	O00215; Q9BW98; A8K7Z0; B2RXG3; B4E1D5; H3BRV0
UniProt Related Accession #	Q99613
GenBank Nucleotide #	NM_001037808.2
NCBI Official Full Name	eukaryotic translation initiation factor 3 subunit C isoform a
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 3 subunit C
NCBI Official Symbol	EIF3C
NCBI Official Synonym Symbols	EIF3CL; EIF3S8; eIF3-p110
UniProt Protein Name	Eukaryotic translation initiation factor 3 subunit C
UniProt Synonym Protein Names	Eukaryotic translation initiation factor 3 subunit 8
Protein Family	Eukaryotic translation initiation factor
UniProt Gene Name	EIF3C
UniProt Synonym Gene Names	eIF3c
UniProt Entry Name	EIF3C_HUMAN
NCBI Protein Information	eukaryotic translation initiation factor 3 subunit C
UniProt Comments for EIF3C	EIF3C: Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is composed of 13 subunits: EIF3A, EIF3B, EIF3C, EIF3D, EIF3E, EIF3F, EIF3G, EIF3H, EIF3I, EIF3J, EIF3K, EIF3L and EIF3M. The eIF-3 complex appears to include 3 stable modules: module A is composed of EIF3A, EIF3B, EIF3G and EIF3I; module B is composed of EIF3F, EIF3H, and EIF3M; and module C is composed of EIF3C, EIF3D, EIF3E, EIF3K and EIF3L. EIF3C of module C binds EIF3B of module A and EIF3H of module B, thereby linking the three modules. EIF3J is a labile subunit that binds to the eIF-3 complex via EIF3B. The eIF-3 complex interacts with RPS6KB1 under conditions of nutrient depletion. Mitogenic stimulation leads to binding and activation of a complex composed of MTOR and RPTOR, leading to phosphorylation and release of RPS6KB1 and binding of EIF4B to eIF-3. Identified in a HCV IRES-mediated translation complex, at least composed of EIF3C, IGF2BP1, RPS3 and HCV RNA-replicon. Belongs to the eIF-3 subunit C family. Protein type: Translation; Translation initiation; RNA-binding Chromosomal Location of Human Ortholog: 16p11.2 Cellular Component: cytosol; eukaryotic translation initiation factor 3 complex Molecular Function: protein binding; translation initiation factor activity; translation initiation factor binding Biological Process: cellular protein metabolic process; formation of translation preinitiation complex; gene expression; regulation of translational initiation; translation; translational initiation
Research Articles on EIF3C	**1.** These findings suggest that eIF3c is overexpressed in human gliomas and essential for their proliferation and survival.

Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/eif3c/7002795