
PLEC Antibody – Rabbit Polyclonal


PRODUCT OVERVIEW

Product Name	PLEC, Polyclonal Antibody
Full Product Name	PLEC Antibody
Product Gene Name	anti-PLEC antibody
Product Synonym Names	Plectin; PCN; PLTN; Hemidesmosomal protein 1; HD1; Plectin-1; PLEC; PLEC1
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Liquid
Purity/Purification	>95%, Protein G purified
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.
Matching Pairs	Unconjugated Antibody: PLEC (MBS7108259) HRP Conjugated Antibody: PLEC (MBS7108260)
Matching Pairs	Unconjugated Antibody: PLEC (MBS7108259) FITC Conjugated Antibody: PLEC (MBS7108261)
Matching Pairs	Unconjugated Antibody: PLEC (MBS7108259) Biotin Conjugated Antibody: PLEC (MBS7108262)
Molecular Weight	514,775 Da
OMIM	Z54367 Genomic DNA
3D Structure	ModBase 3D Structure for Q15149
Immunogen	Recombinant Human Plectin protein (1744-1929AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Product Categories/Family for anti-PLEC antibody	Signal Transduction
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-PLEC antibody	ELISA (EIA), Immunohistochemistry (IHC), Immunofluorescence (IF)
Application Notes for anti-PLEC antibody	IHC: 1:200-1:500 IF: 1:50-1:200
Related Product Information for anti-PLEC antibody	Interlinks intermediate filaments with microtubules and microfilaments and anchors intermediate filaments to desmosomes or hemidesmosomes. Could also bind muscle proteins such as actin to membrane complexes in muscle. May be involved not only in the filaments network, but also in the regulation of their dynamics. Structural component of muscle. Isoform 9 plays a major role in the maintenance of myofiber integrity.


Other Notes	Small volumes of anti-PLEC antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for PLEC purchase	MBS7108259 is an antibody targeting PLEC. Also known as PLEC Antibody. Reactive in Human. Host: Rabbit.



Immunohistochemistry (IHC) of anti-PLEC antibody
IHC image of MBS7108259 diluted at 1:300 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry (IHC) of anti-PLEC antibody
IHC image of MBS7108259 diluted at 1:300 and staining in paraffin-embedded human lung tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry (IHC) of anti-PLEC antibody
Immunofluorescence staining of Hela cells with MBS7108259 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C.The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG (H+L).

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for PLEC. It may not necessarily be applicable to this product.

NCBI GI #	47607492
NCBI GeneID	5339
NCBI Accession #	NP_000436.2
UniProt Primary Accession #	Q15149
UniProt Secondary Accession #	Q15148; Q16640; Q6S376; Q6S377; Q6S378; Q6S379; Q6S380; Q6S381; Q6S382; Q6S383
UniProt Related Accession #	Q15149
GenBank Nucleotide #	NM_000445.4
NCBI Official Full Name	plectin isoform 1c
NCBI Official Synonym Full Names	plectin

NCBI Official Symbol	PLEC
NCBI Official Synonym Symbols	HD1; PCN; EBS1; EBSO; PLTN; EBSMD; EBSND; EBSOG; EBSPA; PLEC1; LGMD2Q; PLEC1b
UniProt Protein Name	Plectin
UniProt Synonym Protein Names	Hemidesmosomal protein 1; HD1; Plectin-1
Protein Family	Non-motile and phage-resistance protein
UniProt Gene Name	PLEC
UniProt Synonym Gene Names	PLEC1; PCN; PLTN; HD1
NCBI Protein Information	plectin
NCBI Summary for PLEC	Plectin is a prominent member of an important family of structurally and in part functionally related proteins, termed plakins or cytolinkers, that are capable of interlinking different elements of the cytoskeleton. Plakins, with their multi-domain structure and enormous size, not only play crucial roles in maintaining cell and tissue integrity and orchestrating dynamic changes in cytoarchitecture and cell shape, but also serve as scaffolding platforms for the assembly, positioning, and regulation of signaling complexes (reviewed in PMID: 9701547, 11854008, and 17499243). Plectin is expressed as several protein isoforms in a wide range of cell types and tissues from a single gene located on chromosome 8 in humans (PMID: 8633055, 8698233). Until 2010, this locus was named plectin 1 (symbol PLEC1 in human; Plec1 in mouse and rat) and the gene product had been referred to as "hemidesmosomal protein 1" or "plectin 1, intermediate filament binding 500kDa". These names were superseded by plectin. The plectin gene locus in mouse on chromosome 15 has been analyzed in detail (PMID: 10556294, 14559777), revealing a genomic exon-intron organization with well over 40 exons spanning over 62 kb and an unusual 5' transcript complexity of plectin isoforms. Eleven exons (1-1j) have been identified that alternatively splice directly into a common exon 2 which is the first exon to encode plectin's highly conserved actin binding domain (ABD). Three additional exons (-1, 0a, and 0) splice into an alternative first coding exon (1c), and two additional exons (2alpha and 3alpha) are optionally spliced within the exons encoding the acting binding domain (exons 2-8). Analysis of the human locus has identified eight of the eleven alternative 5' exons found in mouse and rat (PMID: 14672974); exons 1i, 1j and 1h have not been confirmed in human. Furthermore, isoforms lacking the central rod domain encoded by exon 31 have been detected in mouse (PMID:10556294), rat (PMID: 9177781), and human (PMID: 11441066, 10780662, 20052759). The short alternative amino-terminal sequences encoded by the different first exons direct the targeting of the various isoforms to distinct subcellular locations (PMID: 14559777). As the expression of specific plectin isoforms was found to be dependent on cell type (tissue) and stage of development (PMID: 10556294, 12542521, 17389230) it appears that each cell type (tissue) contains a unique set (proportion and composition) of plectin isoforms, as if custom-made for specific requirements of the particular cells. Concordantly, individual isoforms were found to carry out distinct and specific functions (PMID: 14559777, 12542521, 18541706). In 1996, a number of groups reported that patients suffering from epidermolysis bullosa simplex with muscular dystrophy (EBS-MD) lacked plectin expression in skin and muscle tissues due to defects in the plectin gene (PMID: 8698233, 8941634, 8636409, 8894687, 8696340). Two other subtypes of plectin-related EBS have been described: EBS-pyloric atresia (PA) and EBS-Ogna. For reviews of plectin-related diseases see PMID: 15810881, 19945614. Mutations in the plectin gene related to human diseases should be named based on the position in NM_000445 (variant 1, isoform 1c), unless the mutation is located within one of the other alternative first exons, in which case the position in the respective Reference Sequence should be used. [provided by RefSeq, Aug 2011]
Research Articles on PLEC	**1.** The Structure of the Plakin Domain of Plectin Reveals an Extended Rod-like Shape.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.

Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/plec/7108259