
EGFR Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	EGFR, Polyclonal Antibody
Full Product Name	Anti-EGFR Antibody, Rabbit Polyclonal
Product Gene Name	anti-EGFR antibody
Product Synonym Names	EGFR Antibody, Rabbit PAb, Antigen Affinity Purified; epidermal growth factor receptor; Anti-ERBB Antibody; Anti-ERBB1 Antibody; Anti-ERRP Antibody; Anti-HER1 Antibody; Anti-mENA Antibody; Anti-NISBD2 Antibody; Anti-PIG61 Antibody
Clonality	Polyclonal
Isotype	Rabbit IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	Liquid; 0.2um filtered solution in PBS
Purity/Purification	Protein A & Antigen Affinity
Specificity	Human EGFR
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Molecular Weight	69,228 Da
OMIM	131550
3D Structure	ModBase 3D Structure for P00533
Immunogen	Recombinant Human EGFR Protein (NP_005219; Met1-Lys642). EGFR specific IgG was purified by Human EGFR affinity chromatography.
Product Categories/Family for anti-EGFR antibody	EGFR Signaling Pathway; Cancer Drug Targets Receptor Tyrosine Kinases (RTKs)
Applications Tested/Suitable for anti-EGFR antibody	Western Blot (WB)(FAQ Protocol), ELISA (EIA)(FAQ Protocol)
Application Notes for anti-EGFR antibody	WB: 1:500-1:2000 ELISA: 1:5000-1:10000
Related Product Information for anti-EGFR antibody	As a member of the epidermal growth factor receptor (EGFR) family, EGFR protein is type I transmembrane glycoprotein that binds a subset of EGF family ligands including EGF, amphiregulin, TGF-alpha, betacellulin, etc. EGFR protein plays a crucial role in signaling pathway in the regulation of cell proliferation, survival and differentiation. Binding of a ligand induces EGFR protein homo- or heterodimerization, the subsequent tyrosine autophosphorylation and initiates various down stream pathways (MAPK, PI3K/PKB and STAT). In addition, EGFR signaling also has been shown to exert action on carcinogenesis and disease progression, and thus EGFR protein is proposed as a target for cancer therapy currently.
Product References and Citations for anti-EGFR antibody	Schlessinger. J. (2000. Cel. signalin. b. recepto. tyrosin. kinases. Cel. 103(2). 211-25. Giaccone. G. (2005. HER1/EGFR-targete. agents. predictin. th. futur. fo. patient. wit. unpredictabl. outcome. t. therapy. Ann. Oncol. 16(4). 538-48. Yarden. Y.. e. al. (2001. Untanglin. th. Erb. signallin. network. Nat. Rev. Mol. Cell. Biol. 2(2). 127-37.

Other Notes	Small volumes of anti-EGFR antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for EGFR purchase	MBS8126391 is an antibody targeting EGFR. Also known as Anti-EGFR Antibody, Rabbit Polyclonal. Reactive in Human. Host: Rabbit.



Western Blot (WB) of anti-EGFR antibody
Anti-EGFR rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lysates/proteins at 30ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size:134 kDa

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for EGFR. It may not necessarily be applicable to this product.

NCBI GI #	29725609
NCBI GeneID	1956
NCBI Accession #	NP_005219
UniProt Primary Accession #	P00533
UniProt Secondary Accession #	O00688; O00732; P06268; Q14225; Q68GS5; Q92795; Q9BZS2; Q9GZX1; Q9H2C9; Q9H3C9; Q9UMD7
UniProt Related Accession #	P00533
GenBank Nucleotide #	NM_005228.4
NCBI Official Full Name	epidermal growth factor receptor isoform a
NCBI Official Synonym Full Names	epidermal growth factor receptor
NCBI Official Symbol	EGFR
NCBI Official Synonym Symbols	ERBB; HER1; mENA; ERBB1; PIG61; NISBD2
UniProt Protein Name	Epidermal growth factor receptor
UniProt Synonym Protein Names	Proto-oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1
Protein Family	Epidermal growth factor receptor
UniProt Gene Name	EGFR
UniProt Synonym Gene Names	ERBB; ERBB1; HER1
UniProt Entry Name	EGFR_HUMAN
NCBI Protein Information	epidermal growth factor receptor

NCBI Summary for EGFR	The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. [provided by RefSeq, Jun 2016]
UniProt Comments for EGFR	EGFR: a receptor tyrosine kinase. This is a receptor for epidermal growth factor (EGF) and related growth factors including TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30, and vaccinia virus growth factor. EGFR is involved in the control of cell growth and differentiation. It is a single-pass transmembrane tyrosine kinase. Ligand binding to this receptor results in receptor dimerization, autophosphorylation (in trans), activation of various downstream signaling molecules and lysosomal degradation. It can be phosphorylated and activated by Src. Activated EGFR binds the SH2 domain of phospholipase C-gamma (PLC-gamma), activating PLC-gamma-mediated downstream signaling. Phosphorylated EGFR binds Cbl, leading to its ubiquitination and degradation. Grb2 and SHC bind to phospho-EGFR and are involved in the activation of MAP kinase signaling pathways. Phosphorylation on Ser and Thr residues is thought to represent a mechanism for attenuation of EGFR kinase activity. EGFR is overexpressed in breast, head and neck cancers, correlating with poor survival. Activating somatic mutations are seen in lung cancer, corresponding to the minority of patients with strong responses to the EGFR inhibitor Iressa (gefitinib). Mutations and amplifications are also seen in glioblastoma, and upregulation is seen in colon cancer and neoplasms. In xenografts, inhibitors synergize with cytotoxic drugs in the inhibition of many tumor types. Inhibitors include: Iressa/ZD1839, Erbitux, Tarceva, and lapatinib. Four alternatively spliced isoforms have been described. Protein type: Kinase, protein; Tumor suppressor; EC 2.7.10.1; Protein kinase, tyrosine (receptor); Protein kinase, TK; Membrane protein, integral; TK group; EGFR family Chromosomal Location of Human Ortholog: 7p12 Cellular Component: extracellular space; endoplasmic reticulum membrane; nuclear membrane; cell surface; focal adhesion; basolateral plasma membrane; integral to membrane; lipid raft; Golgi membrane; membrane; perinuclear region of cytoplasm; apical plasma membrane; cytoplasm; plasma membrane; AP-2 adaptor complex; endosome membrane; nucleus; endosome; receptor complex Molecular Function: identical protein binding; epidermal growth factor receptor activity; epidermal growth factor binding; nitric-oxide synthase regulator activity; transmembrane receptor protein tyrosine kinase activity; receptor signaling protein tyrosine kinase activity; protein phosphatase binding; protein kinase binding; integrin binding; actin filament binding; protein binding; enzyme binding; transmembrane receptor activity; MAP kinase kinase kinase activity; protein heterodimerization activity; ubiquitin protein ligase binding; protein-tyrosine kinase activity; double-stranded DNA binding; chromatin binding; glycoprotein binding; ATP binding Biological Process: circadian rhythm; diterpenoid metabolic process; positive regulation of nitric oxide biosynthetic process; activation of MAPKK activity; nerve growth factor receptor signaling pathway; alkanesulfonate metabolic process; protein insertion into membrane; positive regulation of vasodilation; G1/S-specific positive regulation of cyclin-dependent protein kinase activity; positive regulation of MAP kinase activity; positive regulation of fibroblast proliferation; cell-cell adhesion; cell surface receptor linked signal transduction; ovulation cycle; hair follicle development; positive regulation of superoxide release; negative regulation of mitotic cell cycle; positive regulation of DNA repair; fibroblast growth factor receptor signaling pathway; digestive tract morphogenesis; response to osmotic stress; phospholipase C activation; response to hydroxyisoflavone; hydrogen peroxide metabolic process; positive regulation of transcription from RNA polymerase II promoter; response to oxidative stress; regulation of nitric-oxide synthase activity; response to calcium ion; negative regulation of protein catabolic process; positive regulation of epithelial cell proliferation; negative regulation of apoptosis; negative regulation of epidermal growth factor receptor signaling pathway; tongue development; axon guidance; embryonic placenta development; peptidyl-tyrosine phosphorylation; translation; protein amino acid autophosphorylation; positive regulation of smooth muscle cell proliferation; signal transduction; positive regulation of synaptic transmission, glutamatergic; learning and/or memory; salivary gland morphogenesis; positive regulation of cell proliferation; response to stress; regulation of peptidyl-tyrosine phosphorylation; epidermal growth factor receptor signaling pathway; ossification; phosphoinositide-mediated signaling; MAPKKK cascade; liver development; positive regulation of protein kinase B signaling cascade; cell proliferation; cerebral cortex cell migration; calcium-dependent phospholipase A2 activation; positive regulation of vasoconstriction; innate immune response; positive regulation of protein amino acid phosphorylation; astrocyte activation; response to cobalamin; positive regulation of phosphorylation; positive regulation of DNA replication; positive regulation of inflammatory response; lung development; positive regulation of cell migration Disease: Lung Cancer

Research Articles on EGFR	**1.** EGFR gene amplification and mutations are rare in triple-negative breast cancer (TNBC), the latter of no apparent clinical relevance. Surrogate markers of EGFR-related chromosomal aberrations and combined EGFR/p53 IHC phenotypes appear to be associated with favorable prognosis in patients with operable TNBC receiving conventional adjuvant chemotherapy.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/egfr/8126391