HMG-1 Antibody – Polyclonal

PRODUCT OVERVIEW

Product Name	HMG-1, Polyclonal Antibody
Full Product Name	HMG-1 Polyclonal Antibody
Product Gene Name	anti-HMG-1 antibody
Clonality	Polyclonal
Molecular Weight	24,894 Da
OMIM	163905
3D Structure	ModBase 3D Structure for P09429
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Related Product Information for anti-HMG-1 antibody	MBS8526477 is an antibody targeting HMG-1. Also known as HMG-1 Polyclonal Antibody.
Other Notes	Small volumes of anti-HMG-1 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for HMG-1. It may not necessarily be applicable to this product.

NCBI GI #	926459555
NCBI GeneID	3146
NCBI Accession #	NP_001300821.1
UniProt Primary Accession #	P09429
UniProt Secondary Accession #	Q14321; Q5T7C3; Q6IBE1; A5D8W9
UniProt Related Accession #	P09429
GenBank Nucleotide #	NM_001313892.1
NCBI Official Full Name	high mobility group protein B1
NCBI Official Synonym Full Names	high mobility group box 1
NCBI Official Symbol	HMGB1
NCBI Official Synonym Symbols	HMG1; HMG3; HMG-1; SBP-1
UniProt Protein Name	High mobility group protein B1
UniProt Synonym Protein Names	High mobility group protein 1; HMG-1
Protein Family	High mobility group-T protein
UniProt Gene Name	HMGB1
UniProt Synonym Gene Names	HMG1; HMG-1
NCBI Protein Information	high mobility group protein B1

NCBI Summary for HMG-1	This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2015]
UniProt Comments for HMG-1	Multifunctional redox sensitive protein with various roles in different cellular compartments. In the nucleus is one of the major chromatin-associated non-histone proteins and acts as a DNA chaperone involved in replication, transcription, chromatin remodeling, V(D)J recombination, DNA repair and genome stability. Proposed to be an universal biosensor for nucleic acids. Promotes host inflammatory response to sterile and infectious signals and is involved in the coordination and integration of innate and adaptive immune responses. In the cytoplasm functions as sensor and/or chaperone for immunogenic nucleic acids implicating the activation of TLR9-mediated immune responses, and mediates autophagy. Acts as danger associated molecular pattern (DAMP) molecule that amplifies immune responses during tissue injury (PubMed:27362237). Released to the extracellular environment can bind DNA, nucleosomes, IL-1 beta, CXCL12, AGER isoform 2/sRAGE, lipopolysaccharide (LPS) and lipoteichoic acid (LTA), and activates cells through engagement of multiple surface receptors. In the extracellular compartment fully reduced HMGB1 (released by necrosis) acts as a chemokine, disulfide HMGB1 (actively secreted) as a cytokine, and sulfonyl HMGB1 (released from apoptotic cells) promotes immunological tolerance (PubMed:23519706, PubMed:23446148, PubMed:23994764, PubMed:25048472). Has proangiogdenic activity (). May be involved in platelet activation (). Binds to phosphatidylserine and phosphatidylethanolamide (). Bound to RAGE mediates signaling for neuronal outgrowth (). May play a role in accumulation of expanded polyglutamine (polyQ) proteins such as huntingtin (HTT) or TBP (PubMed:23303669, PubMed:25549101).
Research Articles on HMG-1	**1.** Study revealed that HMGB1 acts as a mediator of neurite degeneration through the identification of the pathological signaling pathway in Alzheimer's disease (AD). The signaling pathway is initiated with the extracellular release of HMGB1, followed by activation of the TLR4-MAP kinases, phosphorylation of MARCKS at Ser46, and neurite degeneration via instability of the actin network.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-antibody/hmg-1/8526477