MHC Class II (HLA-DPB1) Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	MHC Class II (HLA-DPB1), Polyclonal Antibody
Full Product Name	MHC Class II Rabbit pAb
Product Gene Name	anti-HLA-DPB1 antibody
Product Synonym Names	HLA class II histocompatibility antigen, DP beta 1 chain; HLADPB1; HLADP1B; HLADM
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human
Form/Format	In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Sodium Azide and 50% Glycerol.
Purity/Purification	Affinity Chromatography
Preparation and Storage	Store at 4 degree C short term. Store at -20 degree C long term. Avoid freeze/thaw cycle.
Molecular Weight	29kDa
OMIM	142858
3D Structure	ModBase 3D Structure for P04440
Immunogen	A synthesized peptide derived from human MHC Class II
Cellular Localization	Cell Membrane, Endoplasmic Reticulum, Endosome, Golgi Apparatus, Lysosome, Membrane, MHC II
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for anti-HLA-DPB1 antibody	Western Blot (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF), Immunoprecipitation (IP)
Application Notes for anti-HLA-DPB1 antibody	WB: 1:1000-1:5000 IHC: 1:100-1:500 ICC/IF: 1:50-1:200 IP: 1:50
Related Product Information for anti-HLA-DPB1 antibody	HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins.
Other Notes	Small volumes of anti-HLA-DPB1 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for HLA-DPB1 purchase	MBS8543212 is an antibody targeting HLA-DPB1. Also known as MHC Class II Rabbit pAb. Reactive in Human. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for HLA-DPB1. It may not necessarily be applicable to this product.

NCBI GI #	122263
NCBI GeneID	3115

NCBI Accession #	P04440.1
UniProt Primary Accession #	P04440
UniProt Related Accession #	P04440
NCBI Official Full Name	HLA class II histocompatibility antigen, DP beta 1 chain
NCBI Official Synonym Full Names	major histocompatibility complex, class II, DP beta 1
NCBI Official Symbol	HLA-DPB1
NCBI Official Synonym Symbols	DPB1; HLA-DP; HLA-DPB; HLA-DP1B
UniProt Protein Name	HLA class II histocompatibility antigen, DP beta 1 chain
UniProt Synonym Protein Names	HLA class II histocompatibility antigen, DP(W4) beta chain; MHC class II antigen DPB1
Protein Family	HLA class II histocompatibility antigen
UniProt Gene Name	HLA-DPB1
UniProt Synonym Gene Names	HLA-DP1B
UniProt Entry Name	DPB1_HUMAN
NCBI Protein Information	HLA class II histocompatibility antigen, DP beta 1 chain
NCBI Summary for HLA-DPB1	HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different molecules. [provided by RefSeq, Jul 2008]

UniProt Comments for HLA-DPB1	HLA-DPB1: Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form a heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal miroenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading. Belongs to the MHC class II family. Protein type: Membrane protein, integral Chromosomal Location of Human Ortholog: 6p21.3 Cellular Component: Golgi membrane; cell surface; membrane; lysosomal membrane; plasma membrane; endosome membrane; trans-Golgi network membrane; MHC class II protein complex Molecular Function: peptide antigen binding Biological Process: positive regulation of interferon-gamma production; cytokine and chemokine mediated signaling pathway; T cell costimulation; antigen processing and presentation of exogenous peptide antigen via MHC class II; positive regulation of T cell proliferation; positive regulation of T cell activation; T cell receptor signaling pathway
Research Articles on HLA-DPB1	**1.** The three alleles HLA-B58:01, HLA-DPB110:01 and HLA-DRB111:01 may protect against HIV-2 disease progression towards AIDS.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.