# TRAP220 Antibody – Rabbit Polyclonal

## PRODUCT OVERVIEW

| | |
|---|---|
| **Product Name** | TRAP220, Polyclonal Antibody |
| **Full Product Name** | TRAP220 Antibody |
| **Product Gene Name** | anti-TRAP220 antibody |
| **Product Synonym Gene Name** | MED1 |
| **Product Synonym Names** | ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; Mediator of RNA polymerase II transcription subunit 1; Activator-recruited cofactor 205 kDa component; ARC205; Mediator complex subunit 1; Peroxisome proliferator-activated receptor-binding protein; PBP; PPAR-binding protein; Thyroid hormone receptor-associated protein complex 220 kDa component; Trap220; Thyroid receptor-interacting protein 2; TR-interacting protein 2; TRIP-2; Vitamin D receptor-interacting protein complex component DRIP205; p53 regulatory protein RB18A |
| **Clonality** | Polyclonal |
| **Isotype** | IgG |
| **Host** | Rabbit |
| **Species Reactivity** | Human, Mouse, Rat, Monkey |
| **Form/Format** | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH7.3, 30% glycerol, and 0.01% sodium azide. |
| **Purity/Purification** | The antibody was purified by immunogen affinity chromatography. |
| **Specificity** | Recognizes endogenous levels of TRAP220 protein. |
| **Preparation and Storage** | Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles. |
| **Molecular Weight** | 168,478 Da |
| **Chromosome Location** | Chromosome: 17; NC_000017.10 (37560538..37607527, complement). Location: 17q12 |
| **OMIM** | 604311 |
| **3D Structure** | ModBase 3D Structure for Q15648 |
| **Immunogen** | KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TRAP220. The exact sequence is proprietary. |
| **Product Categories/Family for anti-TRAP220 antibody** | Primary Antibody |
| **ISO Certification** | Manufactured in an ISO 9001:2015 Certified Laboratory. |
| **Applications Tested/Suitable for anti-TRAP220 antibody** | Western Blot (WB), Immunofluorescence (IF), Immunocytochemistry (ICC) |
| **Application Notes for anti-TRAP220 antibody** | WB: 1:500-1:1000<br>IHC: 1:50-1:200 |
| **Related Product Information for anti-TRAP220 antibody** | MBS8583764 is an antibody targeting TRAP220. Also known as TRAP220 Antibody. Reactive in Human, Mouse, Rat, Monkey. Host: Rabbit. Rabbit polyclonal antibody to TRAP220 |
| **Other Notes** | Small volumes of anti-TRAP220 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply. |

## ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for TRAP220. It may not necessarily be applicable to this product.

| | |
|---|---|
| **NCBI GI #** | 28559039 |
| **NCBI GeneID** | 5469 |
| **NCBI Accession #** | NP_004765.2 |
| **UniProt Primary Accession #** | Q15648 |
| **UniProt Secondary Accession #** | O43810; O75447; Q6P9H7; Q6PK58; Q9HD39; A2RRQ6 |
| **UniProt Related Accession #** | Q15648 |
| **GenBank Nucleotide #** | NM_004774.3 |
| **NCBI Official Full Name** | mediator of RNA polymerase II transcription subunit 1 |
| **NCBI Official Synonym Full Names** | mediator complex subunit 1 |
| **NCBI Official Symbol** | MED1 |
| **NCBI Official Synonym Symbols** | PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220 |
| **UniProt Protein Name** | Mediator of RNA polymerase II transcription subunit 1 |
| **UniProt Synonym Protein Names** | Activator-recruited cofactor 205 kDa component; ARC205; Mediator complex subunit 1; Peroxisome proliferator-activated receptor-binding protein; PBP; PPAR-binding protein; Thyroid hormone receptor-associated protein complex 220 kDa component; Trap220; Thyroid receptor-interacting protein 2; TR-interacting protein 2; TRIP-2; Vitamin D receptor-interacting protein complex component DRIP205; p53 regulatory protein RB18A |
| **Protein Family** | Mediator of RNA polymerase II transcription |
| **UniProt Gene Name** | MED1 |
| **UniProt Synonym Gene Names** | ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; ARC205; PBP; PPAR-binding protein; Trap220; TR-interacting protein 2; TRIP-2 |
| **UniProt Entry Name** | MED1_HUMAN |
| **NCBI Protein Information** | mediator of RNA polymerase II transcription subunit 1; ARC205; TRIP-2; PPAR binding protein; PPAR-binding protein; PPARG binding protein; TR-interacting protein 2; p53 regulatory protein RB18A; thyroid receptor interacting protein 2; thyroid receptor-inte |
| **NCBI Summary for TRAP220** | The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. It also regulates p53-dependent apoptosis and it is essential for adipogenesis. This protein is known to have the ability to self-oligomerize. [provided by RefSeq, Jul 2008] |

| | |
|---|---|
| **UniProt Comments for TRAP220** | MED1: a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. Also is a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. May regulates p53-dependent apoptosis. Is essential for adipogenesis. This protein is known to have the ability to self-oligomerize. Two splice-variant isoforms have been described.<br><br>Protein type: Nuclear receptor co-regulator; Transcription, coactivator/corepressor<br><br>Chromosomal Location of Human Ortholog: 17q12<br><br>Cellular Component: nucleoplasm; membrane; nucleolus; Srb-mediator complex; nucleus; chromatin<br><br>Molecular Function: ligand-dependent nuclear receptor binding; peroxisome proliferator activated receptor binding; retinoic acid receptor binding; chromatin DNA binding; transcription coactivator activity; receptor activity; transcription factor binding; thyroid hormone receptor coactivator activity; protein binding; ligand-dependent nuclear receptor transcription coactivator activity; vitamin D receptor binding; transcription cofactor activity; LBD domain binding; estrogen receptor binding; chromatin binding; thyroid hormone receptor binding; nuclear hormone receptor binding<br><br>Biological Process: fat cell differentiation; lactation; lens development in camera-type eye; embryonic placenta development; erythrocyte development; regulation of cell cycle; positive regulation of transcription, DNA-dependent; positive regulation of estrogen receptor signaling pathway; positive regulation of mammary gland epithelial cell proliferation; cell morphogenesis; enucleate erythrocyte development; negative regulation of transcription from RNA polymerase II promoter; cellular lipid metabolic process; androgen biosynthetic process; embryonic hindlimb morphogenesis; angiogenesis; positive regulation of keratinocyte differentiation; steroid hormone receptor signaling pathway; transcription initiation from RNA polymerase II promoter; regulation of transcription from RNA polymerase I promoter; organ regeneration; positive regulation of protein import into nucleus, translocation; monocyte differentiation; embryonic hemopoiesis; thyroid hormone generation; liver development; mRNA transcription from RNA polymerase II promoter; keratinocyte differentiation; negative regulation of neuron differentiation; androgen receptor signaling pathway; embryonic heart tube development; gene expression; positive regulation of transcription from RNA polymerase II promoter; brain development; negative regulation of apoptosis |
| **Research Articles on TRAP220** | **1.** MED1 is recruited to the HER2 gene and required for its expression. |
| **Precautions** | All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines. |
| **Disclaimer** | While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice.<br><br>It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information. |
| **Product Note** | Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details. |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures.

**MyBioSource** | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-monkey-mouse-rat-antibody/trap220/8583764