CLK2 Antibody (N-Term) Blocking Peptide

PRODUCT OVERVIEW

Product Name	CLK2, Blocking Peptide
Full Product Name	CLK2 Antibody (N-term) Blocking Peptide
Product Gene Name	CLK2 blocking peptide
Product Synonym Names	Dual specificity protein kinase CLK2; CDC-like kinase 2; CLK2
Form/Format	The synthetic peptide was lyophilized with 100% acetonitrile and is supplied as a powder. Reconstitute with 0.1 ml deionized water for a final concentration of 1 mg/ml.
Specificity	The synthetic peptide sequence used to generate the antibody was selected from the N-term region of human CLK2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C.
Antibody/Peptide Pairs	CLK2 peptide (MBS9226971) is used for blocking the activity of CLK2 antibody (MBS9213721)
Molecular Weight	59,962 Da
OMIM	602989
Cellular Location	Isoform 1: Nucleus. Nucleus speckle. Note: Inhibition of phosphorylation at Ser-142 results in accumulation in the nuclear speckle.
Tissue Location	Endothelial cells.
Related Product Information for CLK2 blocking peptide	Dual specificity kinase acting on both serine/threonine and tyrosine-containing substrates. Phosphorylates serine- and arginine-rich (SR) proteins of the spliceosomal complex. May be a constituent of a network of regulatory mechanisms that enable SR proteins to control RNA splicing and can cause redistribution of SR proteins from speckles to a diffuse nucleoplasmic distribution. Acts as a suppressor of hepatic gluconeogenesis and glucose output by repressing PPARGC1A transcriptional activity on gluconeogenic genes via its phosphorylation. Phosphorylates PPP2R5B thereby stimulating the assembly of PP2A phosphatase with the PPP2R5B-AKT1 complex leading to dephosphorylation of AKT1. Phosphorylates: PTPN1, SRSF1 and SRSF3. Regulates the alternative splicing of tissue factor (F3) pre-mRNA in endothelial cells.
Other Notes	Small volumes of CLK2 blocking peptide vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for CLK2 purchase	MBS9226971 is a blocking peptide targeting CLK2. Also known as CLK2 Antibody (N-term) Blocking Peptide.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for CLK2. It may not necessarily be applicable to this product.

NCBI GI #	1705919
NCBI GeneID	1196
NCBI Accession #	P49760.1
UniProt Primary Accession #	P49760
UniProt Secondary Accession #	Q96CQ0; B1AVS9; B5MBX6
UniProt Related Accession #	P49760

NCBI Official Full Name	Dual specificity protein kinase CLK2
NCBI Official Synonym Full Names	CDC like kinase 2
NCBI Official Symbol	CLK2
UniProt Protein Name	Dual specificity protein kinase CLK2
UniProt Synonym Protein Names	CDC-like kinase 2
Protein Family	Dual specificity protein kinase
UniProt Gene Name	CLK2
UniProt Entry Name	CLK2_HUMAN
NCBI Protein Information	dual specificity protein kinase CLK2
NCBI Summary for CLK2	This gene encodes a dual specificity protein kinase that phosphorylates serine/threonine and tyrosine-containing substrates. Activity of this protein regulates serine- and arginine-rich (SR) proteins of the spliceosomal complex, thereby influencing alternative transcript splicing. Chromosomal translocations have been characterized between this locus and the PAFAH1B3 (platelet-activating factor acetylhydrolase 1b, catalytic subunit 3 (29kDa)) gene on chromosome 19, resulting in the production of a fusion protein. Note that this gene is distinct from the TELO2 gene (GeneID:9894), which shares the CLK2 alias, but encodes a protein that is involved in telomere length regulation. There is a pseudogene for this gene on chromosome 7. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014]
UniProt Comments for CLK2	CLK2: Dual specificity kinase acting on both serine/threonine and tyrosine-containing substrates. Phosphorylates serine- and arginine-rich (SR) proteins of the spliceosomal complex. May be a constituent of a network of regulatory mechanisms that enable SR proteins to control RNA splicing and can cause redistribution of SR proteins from speckles to a diffuse nucleoplasmic distribution. Acts as a suppressor of hepatic gluconeogenesis and glucose output by repressing PPARGC1A transcriptional activity on gluconeogenic genes via its phosphorylation. Phosphorylates PPP2R5B thereby stimulating the assembly of PP2A phosphatase with the PPP2R5B-AKT1 complex leading to dephosphorylation of AKT1. Phosphorylates: PTPN1, SRSF1 and SRSF3. Regulates the alternative splicing of tissue factor (F3) pre-mRNA in endothelial cells. Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. Lammer subfamily. 3 isoforms of the human protein are produced by alternative splicing. Protein type: Kinase, protein; EC 2.7.12.1; Protein kinase, dual-specificity (non-receptor); Protein kinase, CMGC; CMGC group; CLK family Chromosomal Location of Human Ortholog: 1q21 Cellular Component: nucleoplasm; nucleus Molecular Function: protein binding; protein serine/threonine kinase activity Biological Process: negative regulation of gluconeogenesis; protein amino acid autophosphorylation; protein amino acid phosphorylation; regulation of RNA splicing; response to ionizing radiation
Research Articles on CLK2	**1.** Loss of CLK2 in luminal breast cancer cells leads to the upregulation of epithelial-to-mesenchymal transition (EMT)-related genes and a switch to mesenchymal splice variants of several genes.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.

Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/blocking-peptide/clk2/9226971