
# Paxillin (PXN) Antibody – Rabbit Polyclonal

## PRODUCT OVERVIEW

| | |
|---|---|
| **Product Name** | Paxillin (PXN), Polyclonal Antibody |
| **Full Product Name** | Paxillin (Phospho-Ser272) Antibody |
| **Product Gene Name** | anti-PXN antibody |
| **Product Synonym Names** | Paxillin; PXN |
| **Clonality** | Polyclonal |
| **Host** | Rabbit |
| **Species Reactivity** | Human |
| **Form/Format** | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| **Purity/Purification** | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi |
| **Concentration** | 1.0 mg/ml (lot specific) |
| **Specificity** | The Antibody detects endogenous levels of Paxillin only when phosphorylated at Ser272 |
| **Preparation and Storage** | Store at -20 degree C/1 year |
| **Molecular Weight** | 46,604 Da |
| **OMIM** | 602505 |
| **3D Structure** | ModBase 3D Structure for P49023 |
| **Immunogen Type** | Peptide-KLH |
| **Immunogen Description** | Peptide sequence around phosphorylation site of Serine 272(M-A-S(p)-L-S) derived from Human Paxillin. |
| **Modification** | Phospho-Ser272 |
| **Target Name** | Paxillin |
| **Product Categories/Family for anti-PXN antibody** | Phospho-specific Ab |
| **Applications Tested/Suitable for anti-PXN antibody** | Western Blot (WB) |
| **Application Notes for anti-PXN antibody** | Western blotting: 1:500~1:1000 |
| **Related Product Information for anti-PXN antibody** | Cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion). |
| **Other Notes** | Small volumes of anti-PXN antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply. |
| **Searchable Terms for PXN purchase** | MBS9407919 is an antibody targeting PXN. Also known as Paxillin (Phospho-Ser272) Antibody. Reactive in Human. Host: Rabbit. |




**Western Blot (WB) of anti-PXN antibody**
Western blot analysis of extracts from HeLa and JK using Paxillin (Phospho-Ser272) Antibody.The lane on the right is treated with the antigen-specific peptide.

## ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for PXN. It may not necessarily be applicable to this product.

| | |
|---|---|
| **NCBI GI #** | 170932514 |
| **NCBI GeneID** | 5829 |
| **NCBI Accession #** | NP_001074324.1 |
| **UniProt Primary Accession #** | P49023 |
| **UniProt Secondary Accession #** | O14970; O14971; O60360; Q5HYA4; B2RAI3; B7ZMB4 |
| **UniProt Related Accession #** | P49023 |
| **GenBank Nucleotide #** | NM_001080855.2 |
| **NCBI Official Full Name** | paxillin isoform 2 |
| **NCBI Official Synonym Full Names** | paxillin |
| **NCBI Official Symbol** | PXN |
| **UniProt Protein Name** | Paxillin |
| **Protein Family** | Paxillin |
| **UniProt Gene Name** | PXN |
| **UniProt Entry Name** | PAXI_HUMAN |
| **NCBI Protein Information** | paxillin |
| **NCBI Summary for PXN** | This gene encodes a cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion). Alternatively spliced transcript variants encoding different isoforms have been described for this gene. These isoforms exhibit different expression pattern, and have different biochemical, as well as physiological properties (PMID:9054445). [provided by RefSeq, Aug 2011] |

| | |
|---|---|
| **UniProt Comments for PXN** | PXN: a multi-domain cytoskeletal protein involved in actin-membrane attachment that localizes primarily to focal adhesion sites to the extracellular matrix. Phosphorylated by focal adhesion kinase (FAK) and is a component of integrin signaling. Its phosphorylation provides docking sites for recruitment of signaling molecules to focal adhesions. Binds in vitro to vinculin as well as to the SH3 domain of c-SRC and, when tyrosine phosphorylated, to the Crk SH2 domain. Interacts with GIT1, NUDT16L1/SDOS, PARVA and TGFB1I1. Component of cytoplasmic complexes, which also contain GIT1, ARHGEF6 and PAK1. Binds DDEF2. Interacts with unphosphorylated ITGA4. Interacts with RNF5.Three alternatively-spliced isoforms have been described. Isoform beta binds to focal adhesion kinase but weakly to vinculin. Isoform gamma binds to vinculin but only weakly to focal adhesion kinase.<br><br>Protein type: Adaptor/scaffold; Motility/polarity/chemotaxis; Cytoskeletal; Cell adhesion<br><br>Chromosomal Location of Human Ortholog: 12q24.31<br><br>Cellular Component: nucleoplasm; microtubule associated complex; focal adhesion; lamellipodium; cytoplasm; plasma membrane; stress fiber; cell cortex; cytosol<br><br>Molecular Function: integrin binding; protein binding; zinc ion binding; beta-catenin binding; BH4 domain binding; protein kinase binding; vinculin binding<br><br>Biological Process: lamellipodium biogenesis; epidermal growth factor receptor signaling pathway; integrin-mediated signaling pathway; focal adhesion formation; peptidyl-tyrosine phosphorylation; activation of MAPK activity; signal complex assembly; cytoskeleton organization and biogenesis; signal transduction; regulation of cell shape; muscle contraction; branching morphogenesis of a tube; transforming growth factor beta receptor signaling pathway; vascular endothelial growth factor receptor signaling pathway; cell adhesion |
| **Research Articles on PXN** | **1.** findings reveal a novel mechanism by which PTEN inhibits the progression of colon cancer by inhibiting paxillin expression downstream of PI3K/AKT/NF-kappaB pathway |
| **Precautions** | All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines. |
| **Disclaimer** | While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice.<br><br>It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information. |
| **Product Note** | Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details. |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures.

**MyBioSource** | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-antibody/paxillin/9407919