LCK Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	LCK, Polyclonal Antibody
Full Product Name	LCK (Phospho-Tyr394) Polyclonal Conjugated Antibody
Product Gene Name	anti-LCK antibody
Product Synonym Names	Lck (Phospho-Tyr394) Conjugated Antibody; LCK; LSK; LSK-T; P56-LCK
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, Mouse, Rat
Form/Format	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Specificity	The antibody detects endogenous level of Lck only when phosphorylated at tyrosine 394.
Preparation and Storage	Store at 4 degree C for 6 months.
Molecular Weight	58,001 Da
Chromosome Location	Chromosome: 1; NC_000001.10 (32716840..32751766). Location: 1p34.3
OMIM	153390
3D Structure	ModBase 3D Structure for P06239
Immunogen	Peptide sequence around phosphorylation site of tyrosine 394 (N-E-Y (p)-T-A) derived from Human Lck.
Conjugation	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Excitation Emission	AF350: 346nm/442nm
Applications Tested/Suitable for anti-LCK antibody	Most Applications
Application Notes for anti-LCK antibody	Suggested Dilution: AF350 conjugated: most applications: 1:50-1:250 AF405 conjugated: most applications: 1:50-1:250 AF488 conjugated: most applications: 1:50-1:250 AF555 conjugated: most applications: 1:50-1:250 AF594 conjugated: most applications: 1:50-1:250 AF647 conjugated: most applications: 1:50-1:250 AF680 conjugated: most applications: 1:50-1:250 AF750 conjugated: most applications: 1:50-1:250 Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1:50-1:1000

Related Product Information for anti-LCK antibody	Tyrosine kinase that plays an essential role for the selection and maturation of developing T-cell in the thymus and in mature T-cell function. Is constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors and plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, and thereby recruits the associated LCK to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosines residues within the immunoreceptor tyrosines-based activation motifs (ITAMs) in the cytoplasmic tails of the TCRgamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. In addition, contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, and upon engagement of the CD2 molecule, LCK undergoes hyperphosphorylation and activation. Also plays a role in the IL2 receptor-linked signaling pathway that controls T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR.
Other Notes	Small volumes of anti-LCK antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for LCK purchase	MBS9437023 is an antibody targeting LCK. Also known as LCK (Phospho-Tyr394) Polyclonal Conjugated Antibody. Reactive in Human, Mouse, Rat. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for LCK. It may not necessarily be applicable to this product.

NCBI GI #	112789548
NCBI GeneID	3932
NCBI Accession #	NP_001036236.1
UniProt Primary Accession #	P06239
UniProt Secondary Accession #	P07100; Q12850; Q13152; Q5TDH8; Q5TDH9; Q7RTZ3; Q96DW4; Q9NYT8; D3DPP8
UniProt Related Accession #	P06239
GenBank Nucleotide #	NM_001042771.1
NCBI Official Full Name	tyrosine-protein kinase Lck
NCBI Official Synonym Full Names	lymphocyte-specific protein tyrosine kinase
NCBI Official Symbol	LCK
NCBI Official Synonym Symbols	LSK; YT16; p56lck; pp58lck
UniProt Protein Name	Tyrosine-protein kinase Lck
UniProt Synonym Protein Names	Leukocyte C-terminal Src kinase; LSK; Lymphocyte cell-specific protein-tyrosine kinase; Protein YT16; Proto-oncogene Lck; T cell-specific protein-tyrosine kinase; p56-LCK
Protein Family	Proto-oncogene tyrosine-protein kinase
UniProt Gene Name	LCK
UniProt Synonym Gene Names	LSK
UniProt Entry Name	LCK_HUMAN
NCBI Protein Information	tyrosine-protein kinase Lck; leukocyte C-terminal Src kinase; p56(LSTRA) protein-tyrosine kinase; t cell-specific protein-tyrosine kinase; proto-oncogene tyrosine-protein kinase LCK; lymphocyte cell-specific protein-tyrosine kinase; T-lymphocyte specific

NCBI Summary for LCK	This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to the plasma membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants, encoding the same protein, have been described. [provided by RefSeq, Jul 2008]
UniProt Comments for LCK	Lck: a tyrosine kinase of the Src family that is crucial to antigen-receptor signaling in lymphocytes. plays an essential role for the selection and maturation of developing T-cell in the thymus and in mature T-cell function. Is constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors and plays a key role in T-cell antigen receptor(TCR)-linked signal transduction pathways. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, and thereby recruits the associated LCK to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosines residues within the immunoreceptor tyrosines-based activation motifs (ITAMs) in the cytoplasmic tails of the TCRgamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. In addition, contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, and upon engagement of the CD2 molecule, LCK undergoes hyperphosphorylation and activation. Also plays a role in the IL2 receptor-linked signaling pathway that controls T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Binds to the cytoplasmic domain of cell surface receptors, such as CD2, CD4, CD5, CD8, CD44, CD45 and CD122. Also binds to effector molecules, such as PI4K, VAV1, RASA1, FYB and to other protein kinases including CDC2, RAF1, ZAP70 and SYK. Binds to phosphatidylinositol 3'-kinase (PI3K) from T-lymphocytes through its SH3 domain and to the tyrosine phosphorylated form of Sam68 through its SH2 domain. Binds to HIV-1 Nef through its SH3 domain. This interaction inhibits its tyrosine-kinase activity. Overexpression in mice leads to thymic tumors. Aberrant expression is seen in T cell leukemias and colon cancer. The leukemic translocation t(1;7)(p34;q34) has breakpoints at the T cell receptor gene and close to the Lck promoters, can cause increased Lck expression, and in one case, point mutations. A mutated Lck has also been seen in a cell line. One patient with aberrant Lck splicing suffered from SCID-like T cell deficiency. Inhibitor: BMS-279700. Three alternatively spliced isoforms of the human proteinhave been described. Protein type: Oncoprotein; EC 2.7.10.2; Protein kinase, tyrosine (non-receptor); Kinase, protein; Protein kinase, TK; TK group; Src family Chromosomal Location of Human Ortholog: 1p34.3 Cellular Component: pericentriolar material; extrinsic to internal side of plasma membrane; endocytic vesicle; plasma membrane; immunological synapse; intercellular junction; cytosol; lipid raft Molecular Function: protein C-terminus binding; CD8 receptor binding; identical protein binding; non-membrane spanning protein tyrosine kinase activity; protein serine/threonine phosphatase activity; protein phosphatase binding; antigen binding; protein kinase binding; ATPase binding; protein binding; protein-tyrosine kinase activity; protein complex binding; phosphoinositide 3-kinase binding; SH2 domain binding; glycoprotein binding; CD4 receptor binding; ATP binding Biological Process: viral reproduction; nerve growth factor receptor signaling pathway; T cell receptor signaling pathway; protein amino acid phosphorylation; B cell receptor signaling pathway; positive regulation of T cell receptor signaling pathway; hemopoiesis; positive regulation of gamma-delta T cell differentiation; aging; T cell differentiation; regulation of lymphocyte activation; caspase activation; response to drug; epidermal growth factor receptor signaling pathway; platelet activation; fibroblast growth factor receptor signaling pathway; phosphoinositide-mediated signaling; cellular zinc ion homeostasis; regulation of defense response to virus by virus; regulation of cell proliferation; dephosphorylation; positive regulation of tyrosine phosphorylation of Stat5 protein; response to hydrogen peroxide; response to zinc ion; response to mechanical stimulus; release of sequestered calcium ion into cytosol; T cell costimulation; innate immune response; positive regulation of T cell activation; blood coagulation; leukocyte migration; transmembrane receptor protein tyrosine kinase signaling pathway Disease: Immunodeficiency 22
Research Articles on LCK	**1.** conformational states regulate clustering in early T cell signaling

Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-mouse-rat-antibody/lck/9437023