HLA-DRA Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	HLA-DRA, Polyclonal Antibody
Full Product Name	HLA-DRA Conjugated Antibody
Product Gene Name	anti-HLA-DRA antibody
Product Synonym Names	MLRW; HLA-DRA1
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, Mouse, Rat
Form/Format	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Specificity	The antibody detects endogenous level of total HLA-DRA protein.
Preparation and Storage	Store at 4 degree C for 6 months.
Molecular Weight	254
OMIM	gene 610424
3D Structure	ModBase 3D Structure for P01903
Immunogen	Recombinant protein of human HLA-DRA.
Conjugation	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Excitation Emission	AF350: 346nm/442nm
Applications Tested/Suitable for anti-HLA-DRA antibody	Most Applications
Application Notes for anti-HLA-DRA antibody	Suggested Dilution: AF350 conjugated: most applications: 1:50-1:250 AF405 conjugated: most applications: 1:50-1:250 AF488 conjugated: most applications: 1:50-1:250 AF555 conjugated: most applications: 1:50-1:250 AF594 conjugated: most applications: 1:50-1:250 AF647 conjugated: most applications: 1:50-1:250 AF680 conjugated: most applications: 1:50-1:250 AF750 conjugated: most applications: 1:50-1:250 Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1:50-1:1000
Related Product Information for anti-HLA-DRA antibody	HLA-DRA is one of the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha and a beta chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. DRA does not have polymorphisms in the peptide binding part and acts as the sole alpha chain for DRB1, DRB3, DRB4 and DRB5.
Other Notes	Small volumes of anti-HLA-DRA antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.

Searchable Terms for HLA-DRA purchase	MBS9443456 is an antibody targeting HLA-DRA. Also known as HLA-DRA Conjugated Antibody. Reactive in Human, Mouse, Rat. Host: Rabbit.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for HLA-DRA. It may not necessarily be applicable to this product.

NCBI GI #	122206
NCBI GeneID	3122
NCBI Accession #	P01903.1
UniProt Primary Accession #	P01903
UniProt Secondary Accession #	Q30160; Q6IAZ1; Q861I2; Q9TP70; A2BET4
UniProt Related Accession #	P01903
NCBI Official Full Name	HLA class II histocompatibility antigen, DR alpha chain
NCBI Official Synonym Full Names	major histocompatibility complex, class II, DR alpha
NCBI Official Symbol	HLA-DRA
NCBI Official Synonym Symbols	MLRW; HLA-DRA1
UniProt Protein Name	HLA class II histocompatibility antigen, DR alpha chain
UniProt Synonym Protein Names	MHC class II antigen DRA
Protein Family	HLA class II histocompatibility antigen
UniProt Gene Name	HLA-DRA
UniProt Synonym Gene Names	HLA-DRA1
UniProt Entry Name	DRA_HUMAN
NCBI Protein Information	HLA class II histocompatibility antigen, DR alpha chain; MHC class II antigen DRA; MHC cell surface glycoprotein; histocompatibility antigen HLA-DR alpha

UniProt Comments for HLA-DRA	HLA-DRA: Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form a heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal miroenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading. Belongs to the MHC class II family. Protein type: Membrane protein, integral Chromosomal Location of Human Ortholog: 6p21.3 Cellular Component: Golgi membrane; cell surface; lysosome; lysosomal membrane; late endosome membrane; integral to plasma membrane; plasma membrane; trans-Golgi network membrane; MHC class II protein complex Molecular Function: MHC class II receptor activity; peptide antigen binding Biological Process: peptide antigen assembly with MHC class II protein complex; cytokine and chemokine mediated signaling pathway; T cell costimulation; antigen processing and presentation of exogenous peptide antigen via MHC class II; immune response; cognition; T cell receptor signaling pathway; antigen processing and presentation of peptide or polysaccharide antigen via MHC class II; polysaccharide assembly with MHC class II protein complex Disease: Hepatitis B Virus, Susceptibility To
Research Articles on HLA-DRA	**1.** The effects of rs11248051 and rs1564282 variants of GAK, and the rs3129882 variant of HLA-DRA, were investigated in Parkinson's disease patients.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-mouse-rat-antibody/hla-dra/9443456