
CD55 Antibody – Rabbit Polyclonal

PRODUCT OVERVIEW

Product Name	CD55, Polyclonal Antibody
Full Product Name	CD55 Rabbit Polyclonal Antibody
Product Gene Name	anti-CD55 antibody
Product Synonym Names	CD55; CR; CROM; DAF; TC
Clonality	Polyclonal
Isotype	IgG
Host	Rabbit
Species Reactivity	Human, Mouse
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purity/Purification	Affinity purification
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.
Molecular Weight	41,400 Da
Chromosome Location	Chromosome: 1; NC_000001.10 (207494817..207534311). Location: 1q32
OMIM	125240
3D Structure	ModBase 3D Structure for P08174
Immunogen	Recombinant fusion protein of human CD55 (NP_000565.1).
Product Categories/Family for anti-CD55 antibody	Primary Antibody
Applications Tested/Suitable for anti-CD55 antibody	Western Blot (WB)
Application Notes for anti-CD55 antibody	WB: 1:500-1:2000
Related Product Information for anti-CD55 antibody	This gene encodes a glycoprotein involved in the regulation of the complement cascade. Binding of the encoded protein to complement proteins accelerates their decay, thereby disrupting the cascade and preventing damage to host cells. Antigens present on this protein constitute the Cromer blood group system (CROM). Alternative splicing results in multiple transcript variants. The predominant transcript variant encodes a membrane-bound protein, but alternatively spliced transcripts may produce soluble proteins.
Other Notes	Small volumes of anti-CD55 antibody vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for CD55 purchase	MBS9464980 is an antibody targeting CD55. Also known as CD55 Rabbit Polyclonal Antibody. Reactive in Human, Mouse. Host: Rabbit.



Western Blot (WB) of anti-CD55 antibody
Western blot analysis of extracts of A-549 cells, using CD55 antibody.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for CD55. It may not necessarily be applicable to this product.

NCBI GI #	10835143
NCBI GeneID	1604
NCBI Accession #	NP_000565.1
UniProt Primary Accession #	P08174
UniProt Secondary Accession #	P09679; P78361; B1AP14; D3DT83; D3DT84
UniProt Related Accession #	P08174
GenBank Nucleotide #	NM_000574.3
NCBI Official Full Name	complement decay-accelerating factor isoform 1 preproprotein
NCBI Official Synonym Full Names	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
NCBI Official Symbol	CD55
NCBI Official Synonym Symbols	CR; TC; DAF; CROM
UniProt Protein Name	Complement decay-accelerating factor
Protein Family	Complement decay-accelerating factor
UniProt Gene Name	CD55
UniProt Synonym Gene Names	CR; DAF
UniProt Entry Name	DAF_HUMAN
NCBI Protein Information	complement decay-accelerating factor; CD55 antigen
NCBI Summary for CD55	This gene encodes a protein involved in the regulation of the complement cascade. The encoded glycoprotein is also known as the decay-accelerating factor (DAF); binding of DAF to complement proteins accelerates their decay, disrupting the cascade and preventing damage to host cells. Antigens present on the DAF glycoprotein constitute the Cromer blood group system (CROM). Two alternatively spliced transcripts encoding different proteins have been identified. The predominant transcript encodes a membrane-bound protein expressed on cells exposed to plasma component proteins but an alternatively spliced transcript produces a soluble protein present at much lower levels. Additional, alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008]

UniProt Comments for CD55	CD55: This protein recognizes C4b and C3b fragments that condense with cell-surface hydroxyl or amino groups when nascent C4b and C3b are locally generated during C4 and c3 activation. Interaction of daf with cell-associated C4b and C3b polypeptides interferes with their ability to catalyze the conversion of C2 and factor B to enzymatically active C2a and Bb and thereby prevents the formation of C4b2a and C3bBb, the amplification convertases of the complement cascade. Belongs to the receptors of complement activation (RCA) family. 2 isoforms of the human protein are produced by alternative splicing. Protein type: Membrane protein, GPI anchor; Cell surface Chromosomal Location of Human Ortholog: 1q32 Cellular Component: cell surface; integral to plasma membrane; extracellular region; plasma membrane; lipid raft Molecular Function: viral receptor activity; protein binding; lipid binding Biological Process: respiratory burst; entry of virus into host cell; elevation of cytosolic calcium ion concentration; negative regulation of complement activation; regulation of complement activation; innate immune response; regulation of lipopolysaccharide-mediated signaling pathway; complement activation, classical pathway Disease: Blood Group, Cromer System
Research Articles on CD55	**1.** Crry deletion from DAF-deficient mouse platelets cause abnormal platelet turnover, leading to compensatory increase in thrombopoiesis.
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.
Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/polyclonal-human-mouse-antibody/cd55/9464980