Phospho-CAD (Thr456) Blocking Peptide – The purity is >90%,tested by HPLC and MS. Purity

PRODUCT OVERVIEW

Product Name	CAD, Blocking Peptide
Full Product Name	Phospho-CAD (Thr456) Blocking Peptide
Product Gene Name	CAD blocking peptide
Product Synonym Names	Aspartate carbamoyltransferase; Aspartate transcarbamylase; CAD protein; CAD trifunctional protein; Carbamoyl phosphate synthetase 2; Carbamoyl-phosphate synthetase 2 aspartate transcarbamylase and dihydroorotase; CPSase ATCase DHOase; Dihydroorotase; Glutamine dependent carbamoyl phosphate synthase; Multifunctional protein CAD
Form/Format	Synthetic peptide was lyophilized with 100% acetonitrile and is supplied as a powder. Reconstitute with 0.1 ml DI water for a final concentration of 10 mg/ml.
Purity/Purification	The purity is >90%,tested by HPLC and MS.
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C.
Antibody/Peptide Pairs	CAD (Thr456) peptide (MBS9616265) is used for blocking the activity of CAD (Thr456) antibody (MBS9612984)
Molecular Weight	242,984 Da
Chromosome Location	Chromosome: 2; NC_000002.11 (27440258..27466654). Location: 2p22-p21
OMIM	114010
Supply Chain Verification	Manufactured in a lab with traceable raw materials manufactured on site. Coordinated product portfolio of antibodies, pairs, conjugates, recombinant proteins, and immunoassay materials available, please inquire.
Notes	For phospho antibody, we provide phospho peptide (0.5mg) and non-phospho peptide (0.5mg)
ISO Certification	Manufactured in an ISO 9001:2015 Certified Laboratory.
Applications Tested/Suitable for CAD blocking peptide	Blocking/Control
Related Product Information for CAD blocking peptide	Blocking peptides are peptides that bind specifically to the target antibody and block antibody binding. These peptide usually contains the epitope recognized by the antibody. Antibodies bound to the blocking peptide no longer bind to the epitope on the target protein. This mechanism is useful when non-specific binding is an issue, for example, in Western blotting (immunoblot) and immunohistochemistry (IHC). By comparing the staining from the blocked antibody versus the antibody alone, one can see which staining is specific; Specific binding will be absent from the western blot or immunostaining performed with the neutralized antibody.
Other Notes	Small volumes of CAD blocking peptide vial(s) may occasionally become entrapped in the seal of the product vial during shipment and storage. If necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container`s cap. Certain products may require to ship with dry ice and additional dry ice fee may apply.
Searchable Terms for CAD purchase	MBS9616265 is a blocking peptide targeting CAD. Also known as Phospho-CAD (Thr456) Blocking Peptide.

ADDITIONAL DETAILS

NCBI/Uniprot data below describe general gene information for CAD. It may not necessarily be applicable to this product.

NCBI GI #	18105007
NCBI GeneID	790
NCBI Accession #	NP_004332.2

UniProt Primary Accession #	P27708
UniProt Secondary Accession #	Q6P0Q0; D6W552
UniProt Related Accession #	P27708
GenBank Nucleotide #	NM_004341.3
NCBI Official Full Name	CAD protein
NCBI Official Synonym Full Names	carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase
NCBI Official Symbol	CAD
UniProt Protein Name	CAD protein
Protein Family	CAD protein
UniProt Gene Name	CAD
UniProt Entry Name	PYR1_HUMAN
NCBI Protein Information	CAD protein; CAD trifunctional protein; multifunctional protein CAD
NCBI Summary for CAD	The de novo synthesis of pyrimidine nucleotides is required for mammalian cells to proliferate. This gene encodes a trifunctional protein which is associated with the enzymatic activities of the first 3 enzymes in the 6-step pathway of pyrimidine biosynthesis: carbamoylphosphate synthetase (CPS II), aspartate transcarbamoylase, and dihydroorotase. This protein is regulated by the mitogen-activated protein kinase (MAPK) cascade, which indicates a direct link between activation of the MAPK cascade and de novo biosynthesis of pyrimidine nucleotides. [provided by RefSeq, Jul 2008]
UniProt Comments for CAD	CAD: a multifunctional enzyme that catalyzes four enzymatic activities of the pyrimidine pathway, including the rate-limiting first step. The four enzymatic activities involved in de novo pyrimidine synthesis are glutamine amidotransferase, carbamoyl-phosphate synthetase, aspartate transcarbamoylase, and dihydroorotase activities. Provides essential precursors for DNA synthesis. Required for the de novo synthesis of pyrimidine nucleotides and cell growth. Phosphorylation by p70S6K promotes CAD oligomerization, stimulating de novo synthesis of pyrimidines and progression through S phase in mammalian cells. Utilizes glutamine, bicarbonate, and aspartic acid to form a pyrimidine ring that is subsequently attached to ribose to generate a pyrimidine. Protein type: EC 6.3.5.5; EC 2.1.3.2; Transferase; Ligase; Amino Acid Metabolism - alanine, aspartate and glutamate; EC 3.5.2.3; Hydrolase; Nucleotide Metabolism - pyrimidine Chromosomal Location of Human Ortholog: 2p22-p21 Cellular Component: nucleoplasm; cell projection; protein complex; nuclear matrix; membrane; cell soma; plasma membrane; terminal button; nucleus; cytosol Molecular Function: carbamoyl-phosphate synthase (ammonia) activity; identical protein binding; enzyme binding; zinc ion binding; carbamoyl-phosphate synthase (glutamine-hydrolyzing) activity; aspartate carbamoyltransferase activity; ATP binding; protein kinase activity; dihydroorotase activity Biological Process: lactation; organ regeneration; arginine biosynthetic process; nucleobase, nucleoside and nucleotide metabolic process; protein amino acid autophosphorylation; heart development; UTP biosynthetic process; peptidyl-threonine phosphorylation; glutamine metabolic process; response to testosterone stimulus; female pregnancy; liver development; response to caffeine; 'de novo' pyrimidine base biosynthetic process; response to cortisol stimulus; pyrimidine base metabolic process; pyrimidine nucleoside biosynthetic process; glutamine catabolic process; drug metabolic process; urea cycle; response to amine stimulus Disease: Congenital Disorder Of Glycosylation, Type Iz
Research Articles on CAD	**1.** preliminary X-ray diffraction analysis of the dihydroorotase domain of human CAD
Precautions	All of MyBioSource's Products are for scientific laboratory research purposes and are not for diagnostic, therapeutics, prophylactic or in vivo use. Through your purchase, you expressly represent and warrant to MyBioSource that you will properly test and use any Products purchased from MyBioSource in accordance with industry standards. MyBioSource and its authorized distributors reserve the right to refuse to process any order where we reasonably believe that the intended use will fall outside of our acceptable guidelines.

Disclaimer	While every efforts were made to ensure the accuracy of the information provided in this datasheet, MyBioSource will not be liable for any omissions or errors contained herein. MyBioSource reserves the right to make changes to this datasheet at any time without prior notice. It is the responsibility of the customer to report product performance issues to MyBioSource within 30 days of receipt of the product. Please visit our Terms & Conditions page for more information.
Product Note	Select online data sheet information is drawn from bioinformatics databases, occasionally resulting in ambiguous or non-relevant product information. It is the responsibility of the customer to review, verify, and evaluate the information to make sure it matches their requirements before purchasing the kit. Our ELISA Kit assays are dynamic research tools and sometimes they may be updated and improved. If the format of this assay is important to you then please request the current manual or contact our technical support team with a presales inquiry before placing an order. We will confirm the current details of the assay. We cannot guarantee the sample manual posted online is the most current manual, it is intended to serve as an example only. Please refer to the instructions For Use provided with the assay kit for precise details.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

MyBioSource | Tel: +1.858.633.0165 | Fax: +1.858.633.0166 | mybiosource.com/contact-us
Online datasheet: https://www.mybiosource.com/blocking-peptide/cad/9616265